

**SELECTIVE CYCLIC PEPTIDES WITH
MELANOCORTIN-4 RECEPTOR (MC4-R) AGONIST ACTIVITY**

Priority to Provisional Application(s) Under 35 U.S.C. § 119(e)

[0001] This application claims priority under 35 U.S.C. § 119(e) of provisional application(s) Serial No. 60/229,184, filed August 30, 2000.

Background of the Invention

[0002] Obesity is widely recognized as a serious health problem for the developed countries, and has reached epidemic status in the United States. More than 50% of the U.S. population is considered overweight, with >25% diagnosed as clinically obese and at considerable risk for heart disease, non-insulin dependent diabetes mellitus (NIDDM), hypertension, and certain cancers. This epidemic presents a significant burden on the health care system as projected obesity treatment costs of more than \$70 billion annually are expected in the U.S. alone. Strategies for treating obesity include reducing food intake or enhancing the expenditure of energy.

[0003] It has been demonstrated that, when injected into the third ventricle of the brain or intraperitoneally, a cyclic heptapeptide analog of α -melanocyte stimulating hormone (α MSH) having melanocortin-4 receptor (MC4-R) agonist activity caused long lasting inhibition of food intake in mice. This effect was reversible when co-administered with a MC4-R antagonist. (Fan, et al., Nature (1997) 385: 165-168) Therefore, agonists of MC4-R activity would be useful in treating or preventing obesity.


[0004] There are five known melanocortin receptors based on sequence homology that ranges from 35-60% homology between family members (Cone, et al., Rec. Prog. Hormone Res. (1996) 51: 287-318), but these receptors differ in their functions. For

[illegible]

[0006] Bednarek, et al. (Peptides (1999) 20: 401-409) and Bednarek, et al. (Biochem. Biophys. Res. Comm. (1999) 261: 209-213) disclose analogs of the cyclic peptide MT-II. They do not disclose any compound of formula I or II described below.


Detailed Description

[0007] This invention provides a compound of the formula:




I


[0008] In compounds of formula I, R^1 and R^{12} together with X and Y form a phenyl ring; or R^1 is hydrogen, $R^2-(NH)_n-C(=O)-NH-$, or




R^{12} is hydrogen; with either X and Y being each C and the bond between X and Y being a double bond, or with X and Y being each CH and the bond between X and Y being a single bond; R^2 is alkyl having from 1 to 5 carbon atoms, alkenyl having from 2 to 5 carbon atoms, or alkynyl having from 2 to 5 carbon atoms; R^{14} is alkyl having from 1 to 5 carbon atoms; and n is 0 or 1, Q is



wherein R^3 , R^4 and R^5 are independently hydrogen, halo, alkyl having from 1 to 4 carbon atoms, hydroxy or alkoxy having from 1 to 4 carbon atoms, wherein when R^4 is not hydrogen, R^3 and R^5 are both hydrogen; R^6 is hydrogen, alkyl having from 1 to 3 carbon atoms, alkoxy having from 1 to 3 carbon atoms, phenoxy, or halo; and R^{11} and R^{13} are each independently hydrogen, alkyl having 3 or 4 carbons, or cycloalkyl having 5 or 6 carbon atoms or R^{11} and R^{13} are both phenyl; R^7 is O or NH; R^8 is hydrogen or methyl. R^9 is




R^{10} is hydrogen or methyl; p is 0 or 1; m is 0, 1, 2, or 3; Z is



R^{17} is hydrogen or lower alkyl, preferably methyl.


[0009] The dotted bond in the compound of formula I is hydrogenated when X and Y are each -CH-. On the other hand, when the dotted bond is present, Y and X taken together with R^1 and R^{12} do not form a phenyl ring, and X and Y are both tetravalent C atoms.

[0010] This invention also provides a compound of the formula:




II


[0011] In compounds of formula II, R^1 is hydrogen, $R^2-(NH)_n-C(=O)-NH-$, or



R^2 is alkyl having from 1 to 5 carbon atoms, alkenyl having from 2 to 5 carbon atoms, or alkynyl having from 2 to 5 carbon atoms; R^{14} is alkyl having from 1 to 5 carbon atoms; and n is 0 or 1, one of R^3 , R^4 , R^5 and R^6 is hydrogen, halo, alkyl having from 1 to 3 carbon atoms, or alkoxy having from 1 to 3 carbon atoms, and the remainder are hydrogen. R^7 is O or NH. R^8 is hydrogen or methyl. R^9 is



R^{10} is hydrogen or methyl; p is 0 or 1; m is 0, 1, 2, or 3; Z is



[0012] The compounds of formulas I and II as well as Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Ala-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2S,3S) beta methyl-Trp-Lys-NH₂ are agonists of the MC4-R. It is known that agonists of MC4-R activity cause reduction of food intake in a mouse model of human obesity. Therefore these compounds are useful in the treatment or prevention of obesity.

[0013] All of the compounds of formulas I and II exemplified below as well as Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Ala-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2S,3S) beta methyl-Trp-Lys-NH₂ were tested for MC4-R agonist activity and MC1-R agonist activity in the *in vitro* assay described below in Biological Activity Example A. All of the tested compounds had an EC50 for MC4-R agonist activity of less than 500 nM, and all exhibited at least 10-fold greater MC4-R agonist activity than MC1-R agonist activity. In contrast, the compound Ac-Nle-Cyclo(Asp-Lys)-Asp-His-(D)Phe-Arg-Trp-Lys-NH₂ exhibited about equal MC1-R and MC4-R agonist activities.


Nomenclature and Abbreviations

[0014] The term "alkyl" means a straight chain or branched alkyl group and the term "lower alkyl" denotes an alkyl group containing from 1 to 6 carbon atoms. The term "alkenyl" means a straight chain or branched alkenyl group. The term "alkynyl" refers to a straight chain or branched alkynyl group.

[0015] The term "alkoxy" means a group of the formula alkyl-O- in which alkyl is as defined above. The term "phenoxy" means a group of the formula phenyl-O- . Except where otherwise specified, "phenyl" refers to an unsubstituted phenyl ring and "phenoxy" refers to an unsubstituted phenoxy group.

[0016] The term "halo" means a group selected from fluoro, chloro, bromo, and iodo.


[0017] Compounds of formula IA are represented as follows:




wherein R^1 , R^3 , R^4 , R^5 , R^7 , R^8 , R^9 , R^{10} , R^{12} , X, Y, Z and p are as above.

[0018] In the compounds of formula IA, R^1 and R^{12} together with X and Y form a phenyl


ring; or R^1 is hydrogen, $R^2-(NH)_n-C(=O)-NH-$, or



[0019] R^{12} is hydrogen, with either X and Y being each C and the bond between X and Y being a double bond, or with X and Y being each CH and the bond between X and Y being a single bond; R^2 is alkyl having from 1 to 5 carbon atoms, alkenyl having from 2 to 5 carbon atoms, or alkynyl having from 2 to 5 carbon atoms; R^{14} is alkyl having from 1 to 5 carbon atoms; and n is 0 or 1. R^3 , R^4 and R^5 are independently hydrogen, halo, alkyl having from 1 to 4 carbon atoms, hydroxy, or alkoxy having from 1 to 4 carbon atoms; wherein when R^4 is not hydrogen, R^3 and R^5 are both hydrogen. R^7 is O or NH. R^8 is hydrogen or methyl. R^9 is




R^{10} is hydrogen or methyl; p is 0 or 1; m is 0, 1, 2, or 3, Z is




R^{17} is hydrogen or lower alkyl, preferably methyl.

[0020] The dotted bond in formula IA can be hydrogenated. When the dotted line is hydrogenated X and Y are both -CH-. On the other hand, when the dotted bond is present, Y and X taken together with R^1 and R^{12} do not form a phenyl ring, X and Y are both the tetravalent C atoms.

[0021] In one embodiment of the compound of formula IA, X and Y are each CH and the bond between X and Y is hydrogenated to a single bond; Z is




R⁷ is O; R¹ is




R² is alkyl having from 1 to 5 carbon atoms; and R¹⁰ and R¹² are both hydrogen. Examples of such compounds include Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Cit-Trp-Lys-NH₂.


[0022] In another embodiment of the compound of formula IA Z is



R⁷ is NH; R¹ is




R² is alkyl; and R¹⁰ and R¹² are both hydrogen. In a more specific embodiment, X and Y are each CH and the bond between X and Y is hydrogenated to a single bond; n is 0; and R⁹ is




[0023] Examples of such compounds include: Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2)Nal-Lys-NH₂; and penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-N-methyl(2)Nal-Lys-NH₂.


[0024] In another more specific embodiment of the compound of formula IA, Z is



R⁷ is NH; R¹ is




R² is alkyl; and R¹⁰ and R¹² are both hydrogen; and R⁹ is



and R¹⁷ is hydrogen or lower alkyl, preferably methyl.


[0025] Such compounds include those in which X and Y are each CH and the bond between X and Y is a single bond; and one of R³, R⁴ and R⁵ is hydrogen, halo or alkyl and the remainder are hydrogen, for example Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-4-MeApc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Glu-Lys)-Glu-Apc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Orn)-Asp-Apc-(D)Phe-Arg-Trp-Orn-NH₂; Penta-cyclo(Asp-Dbr)-Asp-Apc-(D)Phe-Arg-Trp-Dbr-NH₂; Penta-cyclo(Asp-Dpr)-Asp-Apc-(D)Phe-Arg-Trp-Dpr-NH₂; and Ac-cyclo(Asp-Dpr)-Asp-Apc-(D)Phe-Arg-Trp-Dpr-NH₂.

[0026] Another specific embodiment of the compounds of formula IA, are those compounds in which X and Y are each CH and the bond between X and Y is a single bond; and R¹ is



one of R^3 , R^4 and R^5 is alkoxy, and the remainder are hydrogen; and n is 0, for example Penta-cyclo(Asp-Lys)-Asp-4-MeOApc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-4-EtOApc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-4-iPrOApc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-3-MeOApc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-4-OHApc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-4-ClApc-(D)Phe-Arg-Trp-Lys-NH₂.

[0027] Embodiments of compounds of formula IA include those compounds in which each of R^1 , R^3 , R^4 , R^5 , R^8 and R^{10} is hydrogen; R^7 is NH; R^9 is




and R^{17} is hydrogen or lower alkyl preferably methyl;



and p is 0, for example cyclo(succinic acid-Lys)-succinic acid-Apc-(D)Phe-Arg-Trp-Lys-NH₂, cyclo(maleic acid-Lys)-maleic acid-Apc-(D)Phe-Arg-Trp-Lys-NH₂, cyclo(succinic acid-Dpr)-succinic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂, cyclo(maleic acid-Dpr)-maleic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂.

[0028] In another embodiment of the compound of formula IA, R^1 and R^{12} together with X and Y form a phenyl ring. Examples of such compounds include Cyclo(phthalic acid-Lys)-phthalic acid-Apc-(D)Phe-Arg-Trp-Lys-NH₂; Cyclo(phthalic acid-Dpr)-phthalic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂; and Ac-Nle-cyclo(Cys-Cys)-Cys-Apc-(D)Phe-Arg-Trp-Cys-NH₂.


[0029] Compounds of formula IB are represented by the formula:




[0030] In the compounds of formula IB, R¹, is hydrogen,



R² is alkyl having from 1 to 5 carbon atoms, alkenyl having from 2 to 5 carbon atoms, or alkynyl having from 2 to 5 carbon atoms. R¹⁴ is alkyl having from 1 to 5 carbon atoms. N is 0 or 1. R⁶ is hydrogen, alkyl having from 1 to 3 carbons, alkoxy having from 1 to 3 carbons, phenoxy, or halo; R⁷ is O or NH. R⁸ is hydrogen or methyl. R⁹ is




R^{10} is hydrogen or methyl; p is 0 or 1; m is 0, 1, 2, or 3; Z is




R^{17} is hydrogen or lower alkyl, preferably methyl.

[0031] In an embodiment of the compounds of formula IB, i.e., compounds of formula IB1,


Z is



R^7 is NH ; R^1 is



R^2 is alkyl; R^8 and R^{10} are each hydrogen; and R^9 is




and R¹⁷ is as above.

[0032] In a more specific embodiment of such compounds of formula IB1, R⁶ is hydrogen or alkyl. Examples of such compounds include: Penta-cyclo(Asp-Lys)-Asp-Appc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-2-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-2-iPrAppc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-cyclo(Asp-Lys)-Asp-3-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-4-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂.


[0033] In another more specific embodiment of such compounds of formula IB1, R⁶ is halo. Examples of such compounds include Penta-cyclo(Asp-Lys)-Asp-4-ClAppc-(D)Phe-Arg-Trp-Lys-NH₂.

[0034] In another more specific embodiment of such compounds of formula IB1, R⁶ is alkoxy or phenoxy. Examples of such compounds include: Penta-cyclo(Asp-Lys)-Asp-4-PhOAppc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta- (Asp-Lys)-Asp-3-MeO-Appc-(D)Phe-Arg-Trp-Lys- NH₂.


[0035] Compounds of formula IC are represented as follows:




[0036] In the compounds of formula IC, R¹ is hydrogen or



R² is alkyl having from 1 to 5 carbon atoms, alkenyl having from 2 to 5 carbon atoms, or alkynyl having from 2 to 5 carbon atoms. R¹⁴ is alkyl having from 1 to 5 carbon atoms. n is 0 or 1, R¹¹ and R¹³ are each independently hydrogen, alkyl having 3 or 4 carbon atoms, cycloalkyl having 5 or 6 carbon atoms or R¹¹ and R¹³ are both phenyl; R⁷ is O or NH; R⁸ is hydrogen or methyl; R⁹ is




R¹⁰ is hydrogen or methyl; p is 0 or 1; m is 0, 1, 2, or 3; and Z is




R¹⁷ is hydrogen or lower alkyl, preferably methyl.


[0037] In an embodiment of the compound of formula IC, the compound of formula IC1, Z is



R⁷ is NH; R¹ is




R^2 is alkyl; R^8 and R^{10} are each hydrogen; and R^9 is




[0038] In one more specific embodiment of such compounds of formula IC1, one of R^{11} and R^{13} is alkyl or cycloalkyl and the other is hydrogen. Examples of such claims include: Penta-cyclo(Asp-Lys)-Asp-Achc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-Abc-(D) Phe-Arg-Trp-Lys-NH₂.

[0039] In another more specific embodiment of such compounds of formula IC1, one of R^{11} and R^{13} is phenyl and the other is hydrogen or phenyl. Examples of such compounds include Penta-cyclo(Asp-Lys)-Asp-4-Adpc-(D)Phe-Arg-Trp-Lys-NH₂.

[0040] In one embodiment of the compound of formula II, the compound of formula IIA, Z is



R^1 is




R^2 is alkyl;

R^3 , R^4 , R^5 , R^8 and R^{10} are each hydrogen;

R⁶ is hydrogen, halo, alkyl having from 1 to 3 carbon atoms, or alkoxy having from 1 to 3 carbon atoms;

R⁹ is



and R¹⁷ is as above.


[0041] In one embodiment of the compounds of formula II-A, as described in the preceding paragraph, R⁷ is NH. In a more specific embodiment, R⁷ is NH and R⁶ is hydrogen or alkyl. Examples of such compounds include: Penta-cyclo(Asp-Lys)-Asp-(D,L)-Atc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-(Asp-Lys)-Asp-5-Me-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-(Asp-Lys)-Asp-5-Et-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta-(Asp-Lys)-Asp-5-iPr-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂.

[0042] In another specific embodiment of the compound of formula II-A, R⁷ is NH and R⁶ is halo. Examples of such compounds include: Penta-(Asp-Lys)-Asp-5-BrAtc-(D)Phe-Arg-Trp-Lys-NH₂; and Penta-(Asp-Lys)-Asp-5-ClAtc-(D)Phe-Arg-Trp-Lys-NH₂.


[0043] In another specific embodiment of the compound of formula II-A, R⁷ is NH and R⁶ is alkoxy. Penta-(Asp-Lys)-Asp-5-MeO-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-(Asp-Lys)-Asp-5-EtO-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂; Penta-(Asp-Lys)-Asp-5-iPrO-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂.

[0044] Another embodiment of the compounds of formula II are those compounds of formula II in which Z, R¹ to R⁵ and R⁸ to R¹⁰ are as described above, R⁷ is O and R⁶ is halo. Examples of such compounds include: Penta-(Asp-Lys)-Asp-5-BrAtc-(D)Phe-Cit-Trp-Lys-NH₂; and Penta-(Asp-Lys)-Asp-5-ClAtc-(D)Phe-Cit-Trp-Lys-NH₂.

[0045] In another embodiment of the compound of formula II, the compound of formula II-B are those compounds where Z is -S-S- ;



R^3, R^4, R^5, R^8 and R^{10} are each hydrogen; R^6 is hydrogen or halo; R^7 is NH ; and R^9 is



and R^{17} is as above.

[0046] Examples of such compounds of formula II-B include: Ac-Nle-cyclo(Cys-Cys)-Cys-(D,L)Atc-(D)Phe-Arg-Trp-Cys-NH₂; and Penta-cyclo(Cys-Cys)-Cys-5-Br(D,L)Atc-(D)Phe-Arg-Trp-Cys-NH₂.

[0047] This invention also provides the following compounds: Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Ala-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2S,3S) beta methyl-Trp-Lys-NH₂.

[0048] The nomenclature used to define the peptides is that typically used in the art wherein the amino group at the N-terminus appears to the left and the carboxyl group at the C-terminus appears to the right. By natural amino acids is meant one of the naturally occurring amino acids found in proteins, i.e., Gly, Ala, Val, Leu, Ile, Ser, Thr, Lys, Arg, Asp, Asn, Glu, Gln, Cys, Met, Phe, Tyr, Pro, Trp, and His. Where the amino acid has isomeric forms, it is the L form of the amino acid that is represented unless otherwise explicitly indicated.

[0049] The following abbreviations or symbols are used to represent amino acids, protecting groups, solvents, reagents and the like.

<u>Symbol</u>	<u>Meaning</u>
β -Ala	beta-Alanine
(2)-Nal	(2)-Naphthylalanine
Atc	2-Aminotetraline-2-carboxylic acid
5-BrAtc	5-Bromo-2-aminotetraline-2-carboxylic acid
5-ClAtc	5-Chloro-2-aminotetraline-2-carboxylic acid
5-MeOAtc	5-Methoxy-2-aminotetraline-2-carboxylic acid
5-EtOAtc	5-Ethoxy-2-aminotetraline-2-carboxylic acid
5-iPrOAtc	5-Isopropoxy-2-aminotetraline-2-carboxylic acid
5-MeAtc	5-Methyl-2-aminotetraline-2-carboxylic acid
5-EtAtc	5-Ethyl-2-aminotetraline-2-carboxylic acid
5-iPrAtc	5-Isopropyl-2-aminotetraline-2-carboxylic acid
5-DmaAtc	5-Dimethylamino-2-aminotetraline-2-carboxylic acid
DBr	D-2,4-diaminobutanoic acid
DPr	D-2,3-diaminopropionic acid
Sar	Sarcosine (N-methylglycine)
Cit	Citrulline
Apc	1-Amino-4-phenylcyclohexane-1-carboxylic acid
4-HOApc	1-Amino-4-(4-hydroxyphenyl)cyclohexane-1-carboxylic acid
4-MeOApc	1-Amino-4-(4-methoxyphenyl)cyclohexane-1-carboxylic acid
3-MeOApc	1-Amino-4-(4-methoxyphenyl)cyclohexane-1-carboxylic acid
4-EtOApc	1-Amino-4-(4-ethoxyphenyl)cyclohexane-1-carboxylic acid
4-iPrOApc	1-Amino-4-(4-isopropoxyphenyl)cyclohexane-1-carboxylic acid
4-MeApc	1-Amino-4-(4-methylphenyl)cyclohexane-1-carboxylic acid
4-ClApc	1-Amino-4-(4-chlorophenyl)cyclohexane-1-carboxylic acid
Appc	4-Amino-1-phenylpiperidine-4-carboxylic acid
2-MeAppc	4-Amino-1-(2-methylphenyl)piperidine-4-carboxylic acid

2-iProAppc	4-Amino-1-(2-isopropoxyphenyl)piperidine-4-carboxylic acid
3-MeAppc	4-Amino-1-(3-methylphenyl)piperidine-4-carboxylic acid
3-MeOAppc	4-Amino-1-(3-methoxyphenyl)piperidine-4-carboxylic acid
4-MeAppc	4-Amino-1-(4-methylphenyl)piperidine-4-carboxylic acid
4-ClAppc	4-Amino-1-(4-chlorophenyl)piperidine-4-carboxylic acid
4-PhOAppc	4-Amino-1-(4-phenoxyphenyl)piperidine-4-carboxylic acid
Achc	1-Amino-4-cyclohexylcyclohexane-1-carboxylic acid
Adpc	1-Amino-4-diphenylcyclohexane-1-carboxylic acid
Abc	1-Amino-4- <i>tert</i> -butylcyclohexane-1-carboxylic acid
3-Amb	3-Aminomethyl benzoic acid
4-Amb	4-Aminomethyl benzoic acid
2-Aba	2-Aminobenzoic acid
Bu	Butyl
Penta	pentanoyl
Fmoc	9-Fluorenylmethoxycarbonyl
Pmc	2,2,5,7,8-Pentamethylchroman-6-sulfonyl
Trt	Trityl (triphenylmethyl)
CH ₂ Cl ₂	Methylene chloride
CH ₃ CN	Acetonitrile
DMF	Dimethylformamide
DIPEA	N, N-Diisopropylethylamine
TFA	Trifluoroacetic acid
HOBt	N-Hydroxybenzotriazole
DIC	N, N'-Diisopropylcarbodiimide
BOP	Benzotriazol-1-yloxy-tris-(dimethylamino)phosphonium Hexafluorophosphate
PyBroP	Bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium Hexafluorophosphate
FAB-MS	Fast atom bombardment mass spectrometry
ES-MS	Electrospray mass spectrometry

[0054] Common to chemical syntheses of peptides is the protection of reactive side chain groups of the various amino acid moieties with suitable protecting groups, which will prevent a chemical reaction from occurring at that site until the protecting group is ultimately removed. Usually also common is the protection of the alpha amino group of an amino acid or fragment while that entity reacts at the carboxyl group, followed by the selective removal of the alpha amino protecting group and allow a subsequent reaction to take place at that site. While specific protecting groups have been disclosed in regard to the solid phase synthesis method, it should be noted that each amino acid can be protected by an protective group conventionally used for the respective amino acid in solution phase synthesis.

[0055] Alpha amino groups may be protected by a suitable protecting group selected from aromatic urethane-type protecting groups, such as benzyloxycarbonyl (Z) and substituted benzyloxycarbonyl, such as *p*-chlorobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl, *p*-bromobenzyloxycarbonyl, *p*-biphenyl-isopropoxycarbonyl, 9-fluorenylmethoxycarbonyl (Fmoc) and *p*-methoxybenzyloxycarbonyl (Moz); aliphatic urethane-type protecting groups, such as *t*-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropoxycarbonyl, and allyloxycarbonyl. Herein, Fmoc is the most preferred for alpha amino protection.

[0056] Guanidino groups may be protected by a suitable protecting group selected from nitro, *p*-toluenesulfonyl (Tos), Z, pentamethylchromanesulfonyl (Pmc), adamantyloxycarbonyl, and Boc. Pmc is the most preferred for arginine (Arg).

[0057] All solvents, isopropanol (iPrOH), methylene chloride (CH₂Cl₂), dimethylformamide (DMF) and N-methylpyrrolidinone (NMP) were purchased from Fisher or Burdick & Jackson and were used without additional distillation. Trifluoroacetic acid was purchased from Halocarbon or Fluka and used without further purification. Diisopropylcarbodiimide (DIC) and diisopropylethylamine (DIPEA) was purchased from Fluka or Aldrich and used without further purification. Hydroxybenzotriazole (HOBT) dimethylsulfide (DMS) and 1,2-ethanedithiol (EDT) were purchased from Sigma

Chemical Co. and used without further purification. Protected amino acids were generally of the L configuration and were obtained commercially from Bachem, Advanced ChemTech, or Neosystem. Purity of these reagents was confirmed by thin layer chromatography, NMR and melting point prior to use. Benzhydrylamine resin (BHA) was a copolymer of styrene - 1% divinylbenzene (100-200 or 200-400 mesh) obtained from Bachem or Advanced Chemtech. Total nitrogen content of these resins were generally between 0.3 - 1.2 meq/g.

[0058] High performance liquid chromatography (HPLC) was conducted on a LDC apparatus consisting of Constametric I and III pumps, a Gradient Master solvent programmer and mixer, and a Spectromonitor III variable wavelength UV detector. Analytical HPLC was performed in reversed phase mode using Vydac C₁₈ columns (0.4 x 30 cm). Preparative HPLC separations were run on Vydac columns (2 x 25 cm).

[0059] The linear peptides were preferably prepared using solid phase synthesis by the method generally described by Merrifield, [*J. Amer. Chem. Soc.*, **1963**, 85, 2149], although other equivalent chemical synthesis known in the art could be used as previously mentioned. Solid phase synthesis is commenced from the C-terminal end of the peptide by coupling a protected alpha-amino acid to a suitable resin. Such a starting material can be prepared by attaching an alpha-amino-protected amino acid by an ester linkage to a p-benzyloxybenzyl alcohol (Wang) resin, or by an amide bond between an Fmoc-Linker, such as p-[(R, S)- α -[1-(9H-fluoren-9-yl)-methoxyformamido]-2,4-dimethoxybenzyl]-phenoxyacetic acid (Rink linker) to a benzhydrylamine (BHA) resin. Preparation of the hydroxymethyl resin is well known in the art. Fmoc-Linker-BHA resin supports are commercially available and generally used when the desired peptide being synthesized has an unsubstituted amide at the C-terminus.

[0060] Since the compounds of the present invention are cyclic peptides, prepared by the formation of a lactam or a disulfide bond, the precursor linear peptides are assembled so as to place suitable amino acids or mimetics carrying the appropriate side chain residues in positions in the linear peptides that can ultimately be induced to undergo an

intramolecular amide bond or disulfide bond formation. The lactams are formed through the coupling a side chain amino functionality of a C-terminal amino acid residue with a distal carboxylic acid residue, while the disulfide bond is formed through the oxidative coupling of two cysteine residues appropriately incorporated at the C-terminal and at or near the N-terminus of the precursor linear peptide. For example, in the preparation of the lactam peptides, in the precursor linear pentapeptides the N-cap can be used as a template to incorporate a carboxyl residue, e.g. structure X, or in the case of hexapeptides, the peptide is designed such that the N-terminal amino acid residue is chosen from one of the amino acids containing a suitably protected side chain carboxylic acid group, eg aspartic acid, glutamic acid. In the linear heptapeptides, either aspartic acid or glutamic acid are incorporated as the penultimate residue from the N-terminus. In all the precursor linear peptides, i.e., the hepta- hexa- and pentapeptides, the C-terminal residue is chosen from natural or unnatural amino acid bearing a suitably protected basic side chain residue, capable of forming a amide bond when deprotected, for example lysine, ornithine, 2,3-diaminopropanoic acid, 2,4-diaminobutanoic acid. To form a cyclic peptide containing a disulfide bond, where the precursor is a linear hexapeptide, the peptide is designed such that an appropriately S-protected cysteine residue is incorporated as both the C and N-terminal residues, while if the precursor is a linear heptapeptide the appropriately S-protected cysteine residues are incorporated as both the C terminal residue and the penultimate N-terminal residue as in for example X.

[0061] In general, to prepare the linear peptides, the amino acids or mimetic are coupled onto the Fmoc-Linker-BHA resin using the Fmoc protected form of amino acid or mimetic, with 2 - 5 equivalents of amino acid and a suitable coupling reagent. After couplings, the resin may be washed and dried under vacuum. Loading of the amino acid onto the resin may be determined by amino acid analysis of an aliquot of Fmoc-amino acid resin or by determination of Fmoc groups by UV analysis. Any unreacted amino groups may be capped by reacting the resin with acetic anhydride and diisopropylethylamine in methylene chloride.

8	DMF	2 x 30 sec
9	iPrOH	1 x 30 sec
10	DMF	1 x 30 sec
11	CH ₂ Cl ₂	2 x 30 sec

[0065] Solvents for all washings and couplings were measured to volumes of 10 - 20 ml/g resins. Coupling reactions throughout the synthesis were monitored by the Kaiser ninhydrin test to determine extent of completion [Kaiser *et al. Anal. Biochem.* **1970**, *34*, 595-598]. Slow reaction kinetics was observed for Fmoc-Arg (Pmc) and for couplings to secondary amines by sterically hindered acids. Any incomplete coupling reactions were either recoupled with freshly prepared activated amino acid or capped by treating the peptide resin with acetic anhydride as described above. The fully assembled peptide-resins were dried in vacuum for several hours.

[0066] For each compound, the blocking groups were removed and the linear peptide cleaved from the resin by the following procedure. Generally. The peptide-resins are treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 300 μ L anisole, and 9.5 mL trifluoroacetic acid, per gram of resin, at room temperature for 120 min. The resin is filtered off and the filtrates are precipitated in chilled ethyl ether. The precipitates are centrifuged and the ether layer is decanted. The residue is washed with two or three volumes of Et₂O and recentrifuged. If desired, the crude linear peptides are purified by preparative HPLC. The peptides were applied to the columns in a minimum volume of either AcOH/H₂O or 0.1% TFA/H₂O. Gradient elution is generally started at 10% B buffer, 10% -60% B in 90 minutes, (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) at a flow rate of 8 mL/min. UV detection is made at 280 nm. Fractions were collected at 1.0 - 2.5 minute intervals and inspected by analytical HPLC. Fractions judged to be of high purity were pooled and lyophilized.

[0067] To prepare the lactams, the appropriate unpurified linear peptide is dissolved in a suitable inert solvent, for example N-methyl pyrrolidone or DMF, preferably DMF and adjusted to an apparent pH of 8.0 by the addition of a tertiary amine base, for example N-

methy morpholine, and then is treated with an amide bond forming reagent, preferably BOP. The reaction is conveniently carried out at a temperature between 40 °C and 0 °C, preferably at about room temperature. Purification of the crude cyclic peptides is carried out by preparative HPLC. Gradient elution is generally started at 20% B buffer, 20% - 60% B in 90 minutes, (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) at a flow rate of 8 mL/min. UV detection is made at 280 nm. Fractions were collected and monitored by analytical HPLC. Fractions judged to be of high purity were pooled and lyophilized.

[0068] To prepare the cyclic disulfide peptides, the purified HPLC-purified linear peptide containing two appropriately positioned cysteine residues, is dissolved at a fairly high dilution level in an suitable inert solvent mixture, for example aqueous DMSO and the solution is adjusted to pH 8.0 by the careful addition of ammonium hydroxide. Oxygen is then bubbled into the stirred solution. The reaction is conveniently carried out at a temperature between 40 °C and 0 °C, preferably at about room temperature and the progress of the cyclization is monitored by analytical HPLC. After the reaction is judged to be completed, the solution is lyophilized and the crude cyclic peptide is purified by preparative HPLC. Gradient elution is generally started at 20% B buffer, 20% -60% B in 90 minutes, (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) at a flow rate of 8 mL/min. UV detection is made at 280 nm. Fractions were collected and monitored by analytical HPLC. Fractions judged to be of high purity were pooled and lyophilized.

[0069] Purification of the crude peptides is carried out by preparative HPLC. The peptides were applied to the columns in a minimum volume of either AcOH/H₂O or 0.1% TFA/H₂O. Gradient elution is generally started at 10% B buffer, 10% -60% B in 90 minutes, (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) at a flow rate of 8 mL/min. UV detection is made at 280 nm. Fractions were collected at 1.0 - 2.5 minute intervals and inspected by analytical HPLC. Fractions judged to be of high purity were pooled and lyophilized.

[0070] Purity of the final products is checked by analytical HPLC on a reversed phase column as stated above. Purity of all products is judged to be approximately 95 - 99%. All final products were also subjected to fast atom bombardment mass spectrometry (FAB-MS) or electrospray mass spectrometry (ES-MS). All products yielded the expected parent M+H ions within acceptable limits.

[0071] The process for synthesizing the representative compounds may be carried out by a procedure whereby each amino acid in the desired sequence is added one at a time in succession to another amino acid or residue thereof or by a procedure whereby peptide fragments with the desired amino acid sequence are first synthesized conventionally and then condensed to provide the desired peptide.

[0072] Such conventional procedures for synthesizing the novel compounds of the present invention include for example any solid phase peptide synthesis method. In such a method the synthesis of the novel compounds can be carried out by sequentially incorporating the desired amino acid residues one at a time into the growing peptide chain according to the general principles of solid phase methods [Merrifield, R. B., *J. Amer. Chem. Soc.* **1963**, 85, 2149-2154; Barany *et al.*. The peptides, Analysis, Synthesis and Biology, Vol. 2, Gross, E. and Meienhofer, J., Eds. Academic Press 1-284 (1980)].

[0073] Common to chemical syntheses of peptides is the protection of reactive side chain groups of the various amino acid moieties with suitable protecting groups, which will prevent a chemical reaction from occurring at that site until the protecting group is ultimately removed. Usually also common is the protection of the alpha amino group of an amino acid or fragment while that entity reacts at the carboxyl group, followed by the selective removal of the alpha amino protecting group and allow a subsequent reaction to take place at that site. While specific protecting groups have been disclosed in regard to the solid phase synthesis method, it should be noted that each amino acid can be protected by an protective group conventionally used for the respective amino acid in solution phase synthesis.

[0074] Alpha amino groups may be protected by a suitable protecting group selected from aromatic urethane-type protecting groups, such as benzyloxycarbonyl (Z) and substituted benzyloxycarbonyl, such as *p*-chlorobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl, *p*-bromobenzyloxycarbonyl, *p*-biphenyl-isopropoxycarbonyl, 9-fluorenylmethoxycarbonyl (Fmoc) and *p*-methoxybenzyloxycarbonyl (Moz); aliphatic urethane-type protecting groups, such as *t*-butyloxycarbonyl (Boc), diisopropylmethoxycarbonyl, isopropoxycarbonyl, and allyloxycarbonyl. Herein, Fmoc is the most preferred for alpha amino protection.

[0075] Guanidino groups may be protected by a suitable protecting group selected from nitro, *p*-toluenesulfonyl (Tos), Z, pentamethylchromanesulfonyl (Pmc), adamantyloxycarbonyl, and Boc. Pmc is the most preferred for arginine (Arg).

[0076] All solvents, isopropanol (iPrOH), methylene chloride (CH₂Cl₂), dimethylformamide (DMF) and N-methylpyrrolidinone (NMP) were purchased from Fisher or Burdick & Jackson and were used without additional distillation. Trifluoroacetic acid was purchased from Halocarbon or Fluka and used without further purification. Diisopropylcarbodiimide (DIC) and diisopropylethylamine (DIPEA) was purchased from Fluka or Aldrich and used without further purification. Hydroxybenzotriazole (HOBT) dimethylsulfide (DMS) and 1,2-ethanedithiol (EDT) were purchased from Sigma Chemical Co. and used without further purification. Protected amino acids were generally of the L configuration and were obtained commercially from Bachem, Advanced ChemTech, or Neosystem. Purity of these reagents was confirmed by thin layer chromatography, NMR and melting point prior to use. Benzhydrylamine resin (BHA) was a copolymer of styrene - 1% divinylbenzene (100-200 or 200-400 mesh) obtained from Bachem or Advanced Chemtech. Total nitrogen content of these resins were generally between 0.3 - 1.2 meq/g.

[0077] High performance liquid chromatography (HPLC) was conducted on a LDC apparatus consisting of Constametric I and III pumps, a Gradient Master solvent programmer and mixer, and a Spectromonitor III variable wavelength UV detector.

Analytical HPLC was performed in reversed phase mode using Vydac C₁₈ columns (0.4 x 30 cm). Preparative HPLC separations were run on Vydac columns (2 x 25 cm).

[0078] Peptides were preferably prepared using solid phase synthesis by the method generally described by Merrifield, [*J. Amer. Chem. Soc.*, **1963**, 85, 2149], although other equivalent chemical synthesis known in the art could be used as previously mentioned. Solid phase synthesis is commenced from the C-terminal end of the peptide by coupling a protected alpha-amino acid to a suitable resin. Such a starting material can be prepared by attaching an alpha-amino-protected amino acid by an ester linkage to a p-benzyloxybenzyl alcohol (Wang) resin, or by an amide bond between an Fmoc-Linker, such as p-[(R, S)-α-[1-(9H-fluoren-9-yl)-methoxyformamido]-2,4-dimethyloxybenzyl]-phenoxyacetic acid (Rink linker) to a benzhydrylamine (BHA) resin. Preparation of the hydroxymethyl resin is well known in the art. Fmoc-Linker-BHA resin supports are commercially available and generally used when the desired peptide being synthesized has an unsubstituted amide at the C-terminus.

[0079] In general, the amino acids or mimetic are coupled onto the Fmoc-Linker-BHA resin using the Fmoc protected form of amino acid or mimetic, with 2 - 5 equivalents of amino acid and a suitable coupling reagent. After couplings, the resin may be washed and dried under vacuum. Loading of the amino acid onto the resin may be determined by amino acid analysis of an aliquot of Fmoc-amino acid resin or by determination of Fmoc groups by UV analysis. Any unreacted amino groups may be capped by reacting the resin with acetic anhydride and diisopropylethylamine in methylene chloride.

[0080] The resins are carried through several repetitive cycles to add amino acids sequentially. The alpha amino Fmoc protecting groups are removed under basic conditions. Piperidine, piperazine or morpholine (20-40% v/v) in DMF may be used for this purpose. Preferably 40% piperidine in DMF is utilized.

[0081] Following the removal of the alpha amino protecting group, the subsequent protected amino acids are coupled stepwise in the desired order to obtain an intermediate, protected

peptide-resin. The activating reagents used for coupling of the amino acids in the solid phase synthesis of the peptides are well known in the art. For example, appropriate reagents for such syntheses are benzotriazol-1-yloxy-tri- (dimethylamino) phosphonium hexafluorophosphate (BOP), Bromo-tris-pyrrolidino-phosphonium hexafluorophosphate (PyBroP), 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU), and diisopropylcarbodiimide (DIC). Preferred here are HBTU and DIC. Other activating agents as described by Barany and Merrifield [The Peptides, Vol. 2, J. Meienhofer, ed., Academic Press, 1979, pp 1-284] may be utilized. Various reagents such as 1-hydroxybenzotriazole (HOBT), N-hydroxysuccinimide (HOSu) and 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine (HOOBT) may be added to the coupling mixtures in order to optimize the synthetic cycles. Preferred here is HOBT.

[0082] The protocol for a typical synthetic cycle is as follows:

Protocol 1

<u>Step</u>	<u>Reagent</u>	<u>Time</u>
1	DMF	2 x 30 sec
2	40% piperidine/DMF	1 min
3	40% piperidine/DMF	15 min
4	DMF	2 x 30 sec
5	iPrOH	2 x 30 sec
6	DMF	3 x 30 sec
7	coupling	60 min - 18 hours
8	DMF	2 x 30 sec
9	iPrOH	1 x 30 sec
10	DMF	1 x 30 sec
11	CH ₂ Cl ₂	2 x 30 sec

[0083] Solvents for all washings and couplings were measured to volumes of 10 - 20 ml/g resins. Coupling reactions throughout the synthesis were monitored by the Kaiser ninhydrin test to determine extent of completion [Kaiser *et al. Anal. Biochem.* 1970, 34,

595-598]. Slow reaction kinetics was observed for Fmoc-Arg (Pmc) and for couplings to secondary amines by sterically hindered acids. Any incomplete coupling reactions were either recoupled with freshly prepared activated amino acid or capped by treating the peptide resin with acetic anhydride as described above. The fully assembled peptide-resins were dried in vacuum for several hours.

[0084] For each compound, the blocking groups were removed and the peptide cleaved from the resin by the following procedure. The peptide-resins were treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 300 μ L anisole, and 9.5 mL trifluoroacetic acid, per gram of resin, at room temperature for 120 min. The resin is filtered off and the filtrates are precipitated in chilled ethyl ether. The precipitates are centrifuged and the ether layer is decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged. The crude products are dried under vacuum.


Purification of Crude Peptide Preparations

[0085] Purification of the crude peptides was carried out by preparative HPLC. The peptides were applied to the columns in a minimum volume of either AcOH/H₂O or 0.1% TFA/H₂O. Gradient elution was generally started at 10% B buffer, 10% -60% B in 90 minutes, (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) at a flow rate of 8 mL/min. UV detection was made at 280 nm. Fractions were collected at 1.0 - 2.5 minute intervals and inspected by analytical HPLC. Fractions judged to be of high purity were pooled and lyophilized.


[0086] Purity of the final products was checked by analytical HPLC on a reversed phase column as stated above. Purity of all products was judged to be approximately 95 - 99%. All final products were also subjected to fast atom bombardment mass spectrometry (FAB-MS) or electrospray mass spectrometry (ES-MS). All products yielded the expected parent M+H ions within acceptable limits.

[illegible]


Reaction Scheme A




Reaction Scheme B




Reaction Scheme C




In the same manner:



Reaction Scheme D




1. piperidine, DMF
 2. N-acylation




R^2 , R^8 , R^9 , R^{10} , Q and m
 are as previously described.

18a
18b


1. Piperidine, DMF
 2. $R^2-N=C=O$, DIPEA,
 DMF or CH_2Cl_2




Reaction Scheme E



Reaction Scheme F




14a (R⁷=NH, PG = Pmc)
14b (R⁷=O, PG = H)




15a (R⁷=NH, PG = Pmc)
15b (R⁷=O, PG = H)

1. Piperidine, DMF
 2.




28'


1. Piperidine, DMF
 2.



28'




29a (R⁷=NH, PG = Pmc)
29b (R⁷=O, PG = H)




30a (R⁷=NH, PG = Pmc)
30b (R⁷=O, PG = H)

Reaction Scheme G




29a ($R^7=NH$, PG = Pmc)
29b ($R^7=O$, PG = H)

Trifluoroacetic acid,
 (scavengers)




31a ($R^7=NH$, PG = Pmc)
31b ($R^7=O$, PG = H)

BOP




1




30a ($R^7=NH$, PG = Pmc)
30b ($R^7=O$, PG = H)

Trifluoroacetic acid,
 (scavengers)




32a ($R^7=NH$, PG = Pmc)
32b ($R^7=O$, PG = H)

BOP




1

Reaction Scheme H




R², R³, R⁴, R⁵, R⁶, R⁸, R⁹, R¹⁰, Q, p, and m are as previously described

Reaction Scheme I




$R^2, R^3, R^4, R^5, R^6, R^8, R^9, R^{10}, Q, p,$ and m are as previously described

Reaction Scheme J




Reaction Scheme K



m,p,Q,R³, R⁴, R⁵,R⁶, R⁸ and R¹⁰
are as previously described.


Reaction Scheme L



52a ($R^7=NH$, PG = Pmc)

52b ($R^7=O$, PG = H)

1. piperidine, DMF
2. N-acylation




56a ($R^7=NH$, PG = Pmc)

56b ($R^7=O$, PG = H)

R^2 , R^8 , R^9 , R^{10} , Q and m
are as previously described.

52a
52b


1. Piperidine, DMF
2. $R^2-N=C=O$, DIPEA,
DMF or CH_2Cl_2





57a ($R^7=NH$, PG = Pmc)

57b ($R^7=O$, PG = H)


Reaction Scheme M









(a) i. piperidine, DMF ii acylation
 (b) i. piperidine, DMF ii $R^2-N=C=O$










Reaction Scheme N




Reaction Scheme O




Reaction Scheme Q




Reaction Scheme R




Reaction Scheme S




Reaction Scheme T



Reaction Scheme U



Reaction Scheme V



R^6 is as previously described.

[0088] The linear peptides used herein as the penultimate intermediates in the syntheses of the cyclic peptides of the current invention (Structure 1) are prepared by using conventional solid phase peptide synthesis methodology discussed in the previous section. Each cycle consists of two procedures; the initial cleavage of the Fmoc protecting group from the terminal nitrogen in the resin bound chain followed by acylation of the amine function with an Fmoc protected amino acid. The cycle is generally carried out in accordance with the stepwise procedures outlined in Protocol 1. The deprotection is accomplished by using an organic base, for example piperazine, morpholine or piperidine, preferably piperidine in a suitable inert solvent, for example N,N-dimethylformamide (DMF) or N-methylpyrrolidone (NMP). The coupling reaction can be carried out by one of the many conditions developed for amide bond formation, for example O-benzotriazol-1-yl N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) in the presence of an organic base, for example, diisopropylethylamine (DIPEA) in an inert solvent, for example DMF. Alternatively in the present instance, the amide group can be formed using a carbodiimide, for example, diisopropylcarbodiimide (DIC)

along with an activating agent such as 1-hydroxybenzotriazole (HOBT) in a suitable inert solvent such as DMF.

[0089] In Reaction Scheme A, in the first cycle of the preparation of the linear polypeptides precursors for the cyclic peptides of Structure 1 where Z is NHCO, the Fmoc-Linker-BHA Resin represented by Structure 2 is deprotected and condensed with Fmoc-amino acids of Structure 3 to give the resin bound compounds of 4. In this syntheses of the cyclic peptides, it is required that the Fmoc-amino acid 3 contain a key structural requirement which is a suitably protected basic side chain, that when deprotected, can participate in the formation of the intramolecular amide bond. To extend the growing peptide chain, a second cycle incorporates the Fmoc-amino acids 5 to give the compounds of Structure 6. In the third cycle, treatment of the resin linked peptide 6 furnishes the intermediates of Structure 7a where R⁸ is hydrogen. The intermediates of Structure 7b where R⁸ is methyl are synthesized as shown in Scheme C.

[0090] The intermediates of Structure 7b are prepared from the compounds of Structure 7a, as shown in Scheme C. In this procedure, compounds of Structure 7a, prepared by treating compounds of Structure 6 as described in steps 1-5 of Protocol 1, are reacted with an aryl sulfonyl chloride, preferably 2-nitrobenzenesulfonyl chloride 23 to produce the compounds of Structure 24. This reaction is carried out in the presence of a proton acceptor, for example pyridine, triethylamine (TEA) or DIPEA, preferably DIPEA in a suitable inert solvent, preferably DMF. N-methylation of the formed sulfonamide group in the washed resin bound compounds of Structure 24 is accomplished under Mitsunobu conditions to produce the compound of Structure 25. In carrying out this reaction the sulfonamides of Structure 24 are reacted with methanol in the presence of diethyl azodicarboxylate (DEAD) and triphenylphosphine using methanol as solvent. After the reaction is complete, the resin bound N-methylsulfonamide of Structure 25 is washed free of residual reagents and byproducts.

[0091] In the next steps as set forth in Reaction Scheme A the 2-nitrobenzenesulfonyl residue is removed from Structure 25 by reacting 25 with 2-mercaptoethanol and the strong organic base 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) in a suitable solvent,


preferably DMF, to give the resin bound intermediate of Structure 7b. The third cycle of Reaction Scheme A is completed by coupling compounds of either Structures 7a and 7b with Fmoc-Arg(Pmc)-OH (8) or Fmoc-Cit-OH (9) to give the resin bound compounds of Structure 10.

[0092] The next two cycles shown in Reaction Scheme B are carried out by first reacting peptides of Structure 10 with the amino acid Fmoc-(D)-Phe-OH to produce the compound of Structure 11 and then reacting this compound of Structure 11 with either one of the amino acid mimetics of Structure 12 or 13 to incorporate these amino acids into the resin bound peptide to give the resin bound pentapeptides of either Structures 14 and 15, depending upon which amino acid mimetics is used. The incorporation of additional amino acid containing carboxylic acid side chains into the linear pentapeptides suitable to ultimately participate in the formation of the cyclic peptides of this invention is accomplished in two ways

- a. As seen in Reaction Scheme B, a Fmoc amino acid with a appropriately protected acid side chain is incorporated into the resin bound pentapeptides 14 and 15. Thus in cycle 6 (Reaction Scheme B) Fmoc-Asp(OtBu)-OH (16) or Fmoc-Glu(OtBu)-OH (17) is incorporated into the growing peptide chain to give the resin bound hexapeptides of Structures 18 and 19 respectively or alternatively,
- b. The resin bound pentapeptides 14 and 15 are N-capped with a cyclic anhydride of Structure 28' (Reaction Scheme F) for example, maleic anhydride or phthalic anhydride, to give the compounds of Structure 29 and 30, or alternatively,
- c. As seen in Reaction Scheme B, the resin bound hexapeptides 18 and 19 may be further reacted with an additional amino acid to form a heptapeptide before being N-capped. This is accomplished by the incorporation in the usual manner of an amino acid residue, preferably Fmoc-Nle-OH, to furnish 21 and 22.

[0093] N-capping of heptapeptide or hexapeptide produces the terminal amide function group of the compound 1. In this manner, the substituents X, Y, R¹² and R¹ are produced. To N-cap the resin bound hexapeptides (18, 19) or heptapeptides (21, 22), the polypeptide is first treated with piperidine in DMF to remove the Fmoc protecting group and then is reacted with an acylating agent. As shown in Reaction Scheme D to produce

the compound of Structure 1 before the formation of Z where X and Y are CH, and R¹ is



the resin bound polypeptide of Structure 18 is deprotected and N-acylated to give the resin bound amide of Structure 27 or after deprotection is reacted with an isocyanate to form the ureas of Structure 28. The N-acylation is carried out under a variety of methods well known to one skilled in the art. Among the methods used are:

- (i) reaction of the terminal amino functionality with a carboxylic acid R²-CO₂H in a suitable solvent, such as DMF in the presence of HBTU, and an organic base, preferably DIPEA;
- (ii) reaction of the terminal amino functionality with a carboxylic acid chloride R₂-COCl in a suitable solvent, such as dichloromethane in the presence of an organic base, such as pyridine, TEA and DIPEA, preferably DIPEA; or
- (iii) reaction of the terminal amino functionality with a carboxylic acid anhydride of Structure 28' as shown in Reaction Scheme F. This Reaction is carried out in a suitable solvent, such as dichloromethane or DMF in the presence of an organic base, preferably DIPEA.

[0094] The N-capping reaction in Reaction Scheme D where compounds of Structure 18 are converted to ureas of Structure 28 is carried out by reacting the terminal amino group in the compounds of Structure 28 with an isocyanate R²-NCO. This reaction is carried out in a suitable solvent, such as dichloromethane or DMF in the presence an organic base, preferably DIPEA. When the acylation and urea forming reactions are complete, the resin bound products 27 and 28 are washed free of residual reagents and byproducts. By using similar conditions, N-capping of the resin bound polypeptides of Structure 19, 21, and 22 is carried out by fomation of the N-acylated compounds of Structures 33, 35, 37 and the ureas of Structure 34, 36 and 38 (Reaction Scheme E). However, Reaction Scheme E can be modified to provide R¹⁴ groups other than that derived from Nle by utilizing known

amino acids other than the compound of Structure 20 in Reaction Scheme B to produce compounds of Structures 21 and 22.

[0095] Reaction Schemes G and H illustrate the cleavage of remaining protecting groups in the N-capped polypeptides 29,30, 33 - 38 and the concomitant cleavage of the peptides from the solid support. This reaction is carried out by using a strong organic acid, preferably trifluoroacetic acid, optionally in the presence of an inert solvent such as dichloromethane and a trace (1%) of water and optionally in the presence of one or more carbocation scavengers, for example ethanedithiol, dimethyl sulfide, triethylsilane and anisole. The polypeptide cleavage solution is filtered free from the solid support, then is diluted with a suitable solvent, preferably diethyl ether, and the formed solids are collected by filtration. The solid polypeptides of Structures 39-44 produced in Reaction Scheme H may be purified by reversed phase chromatography by using a preparative C18 column.

[0096] With the appropriate functionalities thus now available to form the intramolecular amide bond, the N-capped linear polypeptides are subjected to amide forming reaction conditions well known in the art. Accordingly each in turn of the the linear peptides 31, 32 (Reaction Scheme G) and 39-44 (Reaction Scheme I) is dissolved in an inert solvent, for example DMSO, and by the addition of a tertiary amine base, for example N-methyl morpholine the solution is adjusted to an apparent pH of 8, prior to the addition of an amide forming reagent, for example BOP. The reaction is conveniently carried out at a temperature of between zero degrees and 40 °C, preferably at about room temperature. The reaction is allowed to proceed until it is judged to be complete. Common methods, used by those of average skill, to monitor the progress of a reaction are for example, TLC or analytical HPLC. After removal of the reaction solvents *in vacuo* the crude cyclic peptides of Structure I where Z is NHCO as shown in Reaction Scheme I may be purified by reversed phase chromatography using a preparative C18 column. In this manner Compound of Structure 1 are produced where Z is NHCO Bridge.

[0097] The cyclic disulfide peptides of Structure 1 where Z is S-S are prepared by methodologies, outlined in Reaction Schemes similar to those described above in Reaction Schemes A through H for the preparation of the lactams of Structure 1 where Z is NHCO. The penultimate linear polypeptides are assembled in a like manner, with the exception that amino acids, that contain protected thiol side chain residues, for example Fmoc-Cys(Trt)-OH, are incorporated at the appropriate positions of the growing resin bound polypeptide, preferably in cycle 1 and 6. This preparation of the resin bound linear polypeptides is illustrated in Reaction Schemes J and K. As previously described and shown in Scheme L, the resin bound linear hexapeptide 52, after removal of the Fmoc protecting group group, may be N-capped either by acylation to give 56 or by reaction with an isocyanate to form the urea 57. In the same way the resin bound hexapeptide 53, and the resin bound heptapeptides 54 and 55 are converted to the corresponding N-acylated derivatives 58, 60 and 62 and the urea derivatives 59, 61 and 63 (Reaction Scheme M).

[0098] In Reaction Scheme N, the N-capped resin bound linear hexapeptide 56 is treated with a strong acid, preferably trifluoroacetic acid, optionally in the presence of an inert solvent such as dichloromethane and optionally in the presence of one or more carbocation scavengers, for example ethanedithiol, dimethyl sulfide, triethylsilane and anisole. This causes cleavage of all side chain protecting groups as well as cleavage of the linear peptide from the solid support. The reaction is conveniently carried out at a temperature of between 0 °C and 35 °C, preferably at room temperature. The polypeptide cleavage solution is filtered free from the solid support, then is diluted with a suitable solvent, preferably diethyl ether, and the formed solids are collected by filtration. The solid polypeptide of Structure 64 thus produced optionally may be purified by reversed phase chromatography by using a preparative C18 column. The linear hexapeptide 64 is then treated under oxidative conditions well known to one skilled in the art, that can induce thiols to form a disulfide bond. Accordingly a dilute aqueous solution of 64 is adjusted to pH 8.0 using a moderately weak inorganic base, preferably ammonium hydroxide, and then oxygen was bubbled into the solution until the cyclization was judged to be complete, using standard methods, for example TLC or HPLC. After

removal of the reaction solvents by lyophilization, the crude cyclic peptide (1; Z= S-S) prepared and isolated in this manner, may be purified by reversed phase chromatography by using a preparative C18 column.

[0099] Under similar conditions to those described hereinbefore in connection with Reaction Schemes H and I, the resin bound linear polypeptides 58-63 as shown in Reaction Schemes O and P, are deprotected and cleaved from the solid support to give the linear peptides 65-70 (Reaction Scheme O), which are then oxidatively cyclized, as described above, to give the corresponding compounds of Structure 1 (Reaction Scheme P).

[0100] The Fmoc-amino acids used in the preparation of the peptides described above, as well as the acylating agents and isocyanates used to N-cap the polypeptides are known compounds that are commercially available. The Fmoc-amino acids 12, including its species, the compound of Structure 13, used in Reaction Scheme B, are prepared as described herein by methods that are well known to those of ordinary skill in the practice of organic chemistry. In Scheme Q, the preparation of Fmoc-amino acid species of the compound of Structure 12 from cyclic ketones is outlined. These species which are the Structure 12 and Structure 13, 79 and 80 are prepared in the same way as the species of Structure 75 as outlined in the procedure of Scheme Q. The 4-phenylcyclohexanones of Structure 71 are converted to the hydantoins of Structure 72 by treatment with ammonium carbonate and potassium cyanide. The reaction is conveniently carried out in aqueous ethanol mixture at a temperature of from 50 °C to 90 °C, preferably between 80 °C and 90 °C. Direct hydrolysis of the hydantoins to the amino acids of Structure 73 require a prolonged treatment with strong base, for example with 6N sodium hydroxide solution or with barium hydroxide at reflux temperature. Alternatively, compounds of Structure 72 can be converted to the bis-Boc derivatives of Structure 74. The reaction is carried out using tert-butyl dicarbonate [(Boc)₂O] in an inert solvent, preferably tetrahydrofuran (THF), in the presence of an organic amine base, preferably TEA and a catalyst, 4-dimethylaminopyridine (DMAP) at a temperature of from zero degrees to room temperature, preferably at room temperature. The bis-Boc hydantoins of Structure 74 are readily converted to the amino acids of Structure 73. The reaction is accomplished

using 1N sodium hydroxide in an inert solvent, preferably dimethoxyethane (DME) at from zero degrees to 50 °C, preferably at about room temperature. Protection of the amino functionality with an Fmoc group in a compound of Structure 73 is carried out under a variety of reaction conditions to give the compound of Structure 75 which is the Fmoc-amino acid species of the compound of Structure 12. The reaction may conveniently be performed by treatment of a solution of the amino acid 73 in a mixture of THF or dioxane, preferably dioxane and aqueous sodium carbonate with 9-fluorenylmethoxycarbonylchloroformate (FmocCl) at a temperature of from zero degrees to room temperature, preferably at room temperature. Alternatively, N-(9-fluorenylmethoxycarbonyloxy)succinimide (FmocOSu) is added to a solution of the amino acid 73 in aqueous acetonitrile containing an organic tertiary amine base, preferably TEA. The reaction is run at from zero degrees to room temperature, preferably at room temperature. In another variation of the procedure, DME is evaporated from the hydrolysis mixture in the conversion of 74 to 73 and the reaction is adjusted to ~pH 11. The resulting solution of the sodium salt of 73 is then treated *in situ* with FmocOSu or FmocCl in dioxane at a temperature of from zero degrees to room temperature, preferably at room temperature. In the same manner, as in Reaction scheme Q, the tetralones 76, the N-aryl-4-ketopiperidines 77, and the cyclohexanone derivatives 78 can be converted to the corresponding Fmoc-amino acids of Structures 13, 79 and 80, all of which along with 73 form subgenera of Structure 12 utilized in Reaction Scheme B.

[0101] In Reaction Scheme Q, the compounds of Structure 73 where R⁴ represents a linear or branched lower alkoxy and both R² and R³ is hydrogen in the sub genus Structure 82, may be prepared by O-alkylation of the compound of Structure 81 as shown in Reaction Scheme R. Where R¹⁶ represents an unbranched lower alkyl moiety, the alkylation is carried out by using a primary alkyl halide of Structure R¹⁶halide in the presence of an alkali metal carbonate, for example, sodium or potassium carbonate. The alkyl halide may be a chloro, bromo or iodo derivative, preferably an alkyl iodide. The reaction may be conveniently carried out in an inert solvent that promotes SN₂ displacement reactions, for example acetone, 2-butanone or N,N-dimethylformamide, preferably acetone, at a temperature of from room temperature to the reflux temperature of the solution,

preferably the reflux temperature. When R¹⁶ represents a branched lower alkyl group, e.g., 2-propyl, the alkylation of the compound of Structure 81 to produce the compound of Structure 82 is carried out by using a secondary alkyl halide of Structure R¹⁶halide in the presence of an alkali metal carbonate, e.g., potassium carbonate. The secondary alkyl halide is preferably a secondary alkyl iodide, for example, 2-iodopropane. The reaction may be conveniently carried out in an inert solvent, preferably N,N-dimethylformamide, at a temperature of from room temperature to the reflux temperature of the solution, preferably at about 100 °C.

[0102] 4-Arylcyclohexanones of Structure 71 which are the starting materials in Reaction Scheme Q can be prepared by methods that are well known to one of ordinary skill in the practice of organic chemistry. As outlined in Scheme S, treatment of the aryl halides of Structure 83, where X¹ represents bromo or iodo with an alkyl metal reagent, preferably t-butyl lithium, results in a transmetalation reaction to give the corresponding aryl lithium of Structure 84. The reaction is conveniently carried out at -78 °C by the addition of a solution of the alkyl lithium into a solution of a compound of Structure 83 in an inert anhydrous solvent, such as diethyl ether or tetrahydrofuran, preferably tetrahydrofuran. The aryl lithium of Structure 84 produced in this manner, is then reacted *in situ* with a solution of the monoketal of cyclohexane-1,4-dione (85) in a suitable inert solvent, for example tetrahydrofuran, while the reaction temperature is maintained below -60 °C, preferably at about -78 °C to give the carbinols of Structure 86. The compounds of Structure 87 are obtained by the dehydration of the carbinols of Structure 86. The reaction is conveniently carried out using a strong organic acid catalyst, preferably p-toluenesulfonic acid in an inert solvent, for example benzene or toluene, preferably benzene, at the reflux temperature of the solvent. The formed water is removed from the reaction mixture by means of a Dean Stark apparatus to enable the reaction to go to completion. Compounds of Structure 88 are produced by hydrogenation of the olefins of Structure 87. The reaction is conveniently carried out using a noble metal catalyst, for example palladium on carbon, in a hydrogen atmosphere in an inert solvent, for example ethanol or ethyl acetate. The hydrogenation is usually carried out at room temperature and 40 psi of hydrogen, however if the aryl ring in Structure 87 contains a group prone to

hydrogenolysis, e.g., if R³, R⁴ or R⁵ represents chloro, the reaction pressure is kept at about 5 psi. Compounds of Structure 88 may be also obtained directly from carbinols of Structure 86 by reductive elimination of the hydroxyl group. In this reaction a solution of the compound of Structure 86 (R³ = R⁴ = H and R⁵ = OMe) in an inert solvent, for example dichloromethane, is treated with a Lewis acid, such as boron trifluoride etherate, and a reducing agent, for example triethylsilane, at a temperature of from zero degrees to room temperature. Removal of the ketal protecting group in compounds of Structure 88 gives the ketone of formula 71, which is the starting material for Reaction Scheme Q for preparing the Fmoc-amino acid species of Structure 75 of the compound of Structure 12. The reaction is conveniently carried out in acetone or 2-butanone, preferably acetone under acid catalysis, for example 4N hydrochloric acid or p-toluenesulfonic acid at from room temperature to the reflux temperature of the reaction mixture, preferably at the reflux temperature.

[0103] 5-Substituted-beta-tetralones of Structure 76 shown in Reaction Scheme Q which is the starting material for preparing compound of Structure 13 are known compounds, or if they are not known they can be prepared by methods that are well known to one of ordinary skill in the field of organic chemistry. In the present instance, compounds of Structure 76 are basically prepared by two methods outlined in Reaction Schemes T and U. As shown in Scheme T, a 2-substituted hydrocinnamic acid of Structure 90 is converted to the corresponding carboxylic acid chloride of Structure 91. This conversion can be carried out by several methods, for example by treatment of the hydrocinnamic acid with oxalyl chloride, optionally in the presence of a catalytic amount of N,N-dimethylformamide, in an inert solvent, such as benzene or dichloromethane, preferably dichloromethane. The reaction may be conveniently carried out at a temperature of from zero degrees to room temperature, preferably at room temperature. Alternatively the compound of Structure 90 is reacted with an acyl chloride forming reagent such as sulfonyl chloride in an inert solvent, for example benzene or toluene, preferably toluene at a temperature between room temperature to the reflux temperature of the solution, preferably at the reflux temperature. The diazoketone of Structure 92 is prepared by treatment of the thus formed acyl halide of Structure 91 in an inert solvent, e.g.,

dichloromethane with an excess of a freshly prepared ethereal solution of diazomethane. The combination of reagents is conveniently carried out at ice bath temperature and the reaction is then allowed to proceed at a temperature of from zero degrees to room temperature, preferably at room temperature. As shown in Reaction Scheme T, cyclization of the diazoketone of Structure 92 to furnish the tetralone of Structure 76 is promoted by rhodium (II) acetate dimer in an inert solvent, e.g., dichloromethane. The reaction is normally carried out at from room temperature to the reflux temperature of the solution, preferably at the reflux temperature.


[0104] Compounds of Structure 76, which is a starting material in Reaction Scheme Q, wherein R⁶ represents a linear or branched lower alkoxy group can be prepared as shown in Reaction Scheme U from compounds of Structure 93. In Reaction Scheme U, the compounds of Structure 94 where R^{15'} is an unbranched lower alkyl moiety are prepared by per-O-alkylation of the naphthalenediol of Structure 93 with a primary alkyl iodide or bromide, preferably an iodide, in the presence of a base such as an alkali metal carbonate, for example, sodium or potassium carbonate. The reaction may be carried out in an inert solvent, preferably N,N-dimethylformamide at a temperature of from room temperature to 100 °C, preferably at 35 °C. The compounds of Structure 97 where R^{15''} is a branched lower alkyl are prepared in two steps from the 2-tetralone of Structure 94. The tetralone of Structure 95 is subjected to dehydrogenation in the presence of a noble metal catalyst, such as palladium metal (10% on carbon) in a suitable high boiling solvent such as p-cymene to give the aromatized compound of Structure 96. The naphthol of Structure 96 is then O-alkylated with a secondary alkyl iodide in the presence of a base such as an alkali metal carbonate, preferably cesium carbonate to furnish the compound of Structure 97. The reaction may be conveniently carried out in an inert solvent, preferably N,N-dimethylformamide at a temperature of from room temperature to 100 °C, preferably at about 40 °C.

[0105] The tetralones of Structures 76 are produced by reduction of the compounds of Structures 94 and 97 under dissolving metal conditions, followed by the acid catalyzed hydrolysis of the intermediate enol ethers. The transformation is conveniently carried out


by the portionwise addition of a large excess of an alkali metal, such as sodium or potassium, preferably sodium, to a boiling solution of the substrate in an lower alcohol, preferably ethanol until the starting material is consumed. The tetralones of Structures 76 are obtained by treatment of a solution of the isolated intermediate enol ethers with a strong acid catalyst, preferably p-toluenesulfonic acid. The hydrolysis may be conveniently carried out in a mixture of a lower alcohol, preferably ethanol, and water at a temperature of between room temperature and the reflux temperature of the solution, preferably at the reflux temperature.

[0106] In preparing the compounds of Structure 77, utilized in Reaction Scheme Q as starting materials in producing the compounds of formula 79 which are the Fmoc-amino acid species of the compound of Structure 12, the compound of Structure 98 is used as a starting material. This is shown in Reaction Scheme V. In Reaction Scheme V, compounds of Structure 100 can be prepared by reactions that are known *per se*. For example, they can be prepared by coupling the secondary amine of Structure 98 with an aryl bromide or iodide, preferably an aryl iodide of Structure 99 (Reaction Scheme V). The coupling reaction is catalyzed by a noble metal catalyst, preferably tri(dibenzylideneacetone)dipalladium, in the presence of a chelating phosphine ligand, preferably tri-o-tolylphosphine, and a hindered alkoxide base such as sodium tert-butoxide. The reaction is conveniently carried out in an inert atmosphere using an anhydrous solvent such as dioxane or toluene, preferably dioxane, at a temperature of from 60 °C to the reflux temperature, preferably at 90 °C. Removal of the carbonyl protecting group in compound 100 to give compounds of Structure 77 can be carried out by a variety of methods well known in the field of organic chemistry. For example, the deprotection can be achieved by treatment of a solution of compound 100 in a low boiling ketone such as acetone or 2-butanone with an aqueous mineral acid solution, for example 6N hydrochloric acid at a temperature of from room temperature to the reflux temperature of the mixture, preferably at the reflux temperature. The compound of Structure 100 when treated in this manner with an aqueous mineral acid forms, in Reaction Scheme V, the compound of Structure 77. The compound of Structure 77, as demonstrated in the discussion of Reaction Scheme Q is the intermediate for the


compound of Structure 79 which is the species of acid of Structure 12 wherein Q is



[0107] In Reaction Scheme A, amino acids of Structure 5 are known compounds where R⁹ is




On the other hand, amino acids of Structure 5 where R¹⁷ in R⁹ is lower alkyl, i.e., where R⁹ is



are prepared from a compound of formula 101 via Reaction Scheme V as follows:

Reaction Scheme W



[0108] As shown in Reaction Scheme W, β -methyl(Nin-Mes)tryptophan **109** was prepared by using the method previously described by Boteju, L.W., Wenger K., and Hruby, V.J. *Tet.Lett.*, **33**, 7491 (1992). In the first step the nitrogen in *trans*-indole-3-acrylic acid **101** is protected by conversion to the corresponding mesitylenesulfonamide **102**. The reaction

is carried out by treating the indole acid 101 with a excess (> 2 equiv.) of a solution of an alkyl or aryl lithium reagent, for example phenyl lithium or n-butyllithium, preferably n-butyllithium in an inert solvent, preferably tetrahydrofuran, at a temperature of from -40 °C to about -100 °C, most conveniently at -78 °C. While the reaction is maintained at about -78 °C, the formed dilithiated species is then reacted with mesitylenesulfonyl chloride to furnish the mesitylenesulfonamide 102. The N-protected indole acrylic acid 102 is then coupled with the chiral auxiliary (R)-4-phenyl-2-oxazolidinone (preparation see Nicolas et al., *J. Org Chem.* **1993**, *58*, 766-770.) as its N-lithiated species 104 to give the chiral acrylamide 105. The coupling was accomplished via a mixed anhydride formed from 102. To form the mixed anhydride 103, the N-protected indole acrylic acid 102 was reacted with a suitable acyl chloride, for example *t*-butyl chloroformate, 2,4,6 trichlorobenzoyl chloride or pivaloyl chloride, preferably pivaloyl chloride in the presence of an tertiary amine base, for example triethylamine or diisopropylethylamine, preferably triethylamine. The criteria that make possible the choice of a suitable acyl chloride to form the anhydride 103 are well established and are known by one of ordinary skill in organic chemistry. The anhydride formation is run in an inert solvent, for example tetrahydrofuran at an initial temperature of between -100 °C and 0 °C, preferably at about -78 °C. The reaction is allowed to go to completion at a temperature of between -78 °C and 0 °C degrees, preferably at about 0 °C. The thus formed mixed anhydride 103 is then reacted *in situ* with a solution of N-lithiated (R)-4-phenyl-2-oxazolidinone 104, previously prepared by treating a solution of (R)-4-phenyl-2-oxazolidinone in an inert solvent, for example tetrahydrofuran, with an equimolar amount of a solution of an alkyl or aryl lithium reagent, for example phenyl lithium or n-butyllithium, preferably n-butyllithium in an inert solvent, for example, hexane at a temperature of between -100 °C and zero degrees, preferably at about -78 °C. The coupling reaction which yields the chiral acrylamide 105 is carried out at an initial temperature of between -100 °C and zero degrees, preferably at about -78 °C, and after all the reagents are combined the reaction is allowed to proceed at a temperature of between -78 °C and room temperature, preferably at about room temperature.

[0109] The highlight of the transformation of the chiral acrylamide 105 into 106, is the controlled generation of the two new contiguous chiral centers present in 106. The reaction involves the stereoselective 1,4- conjugate addition (Michael addition) of a methyl cuprate, formed *in situ* from cuprous bromide-dimethylsulfide complex and methyl magnesium bromide, to the Micheal acceptor, the α,β -unsaturated carbonyl system present in 105. The resultant metal-chelated enolate is then directly halogenated with a halogenating agent, preferably N-bromosuccinimide, to give 106. Again, as in the case of the Michael addition, the incorporation of the bromo atom is stereoselective, controlled by the bulky phenyl group on the chiral auxiliary which effectively shields the *si* face of both the α,β -unsaturated acyloxazolidinone system and the intermediate metal-chelated enolate from attack from incoming reagents. To prepare the methyl cuprate, a solution of methyl magnesium bromide in diethyl ether is added to a solution of cuprous bromide-dimethylsulfide (1:1) complex in an inert solvent, for example dimethyl sulfide or tetrahydrofuran, preferably a mixture thereof. The reaction is run at a temperature of between -78°C and room temperature, preferably at -4°C . Instead of methyl magnesium bromide, any lower alkyl bromide can be added to form R^{17} where R^{17} is a lower alkyl group other than methyl. To this formed solution of methyl cuprate, is added *in situ* a solution of the α,β -unsaturated acyloxazolidinone 105 in an inert solvent, preferably tetrahydrofuran. The methyl cuprate addition is run initially at a temperature of between -30°C and room temperature, preferably at -4°C and then is allowed to proceed at room temperature. When it is judged that the reaction is completed (e.g. analysis by TLC or HPLC), it is then cooled to a temperature of between -100°C and -40°C , preferably at about -78°C , whereupon a solution of a halogenating agent, preferably N-bromosuccinimide in an inert solvent, for example, tetrahydrofuran is added. The reaction is then allowed to proceed at a temperature of between 0°C and room temperature, preferably at about room temperature to give after isolation, the bromide 106. The bromide is the displaced by azide ion with a concomitant inversion of configuration. This transformation is accomplished by reaction of the bromide 106 with tetra-butylammonium azide in the presence of excess sodium azide in an inert solvent, for example acetonitrile to give the azide 107. The reaction is conveniently run at a temperature of between 80°C and room temperature, preferably at about room

temperature. Treatment of 107 and an alkali metal hydroxide, for example sodium, potassium or lithium hydroxide, preferably lithium hydroxide in the presence of hydrogen peroxide resulted in the hydrolysis of the chiral auxiliary to give the azido acid 108. The hydrolysis reaction is run in an inert solvent, preferably water, at a temperature of between 0 °C and room temperature, preferably at about 0 °C. Hydrogenation of the α -azido acid 108 gives the β -methyl(Nin-Mes)tryptophan which is immediately converted to the corresponding N(α)-Fmoc- β -methyl(Nin-Mes)tryptophan 109. The hydrogenation of 108 is carried out over a noble metal catalyst, preferably 10% Pd/C, in an inert solvent, for example a lower alkanol, preferably methanol at low pressure (< 2 atmos) and at room temperature. After removal of the catalyst by filtration, the volatiles are removed and the product is dissolved in an inert solvent, for example tetrahydrofuran or water, preferably a mixture thereof, and treated with a mild inorganic base, for example an alkali metal bicarbonate, preferably sodium bicarbonate, and a Fmoc N-protecting group forming reagent, for example 9-fluorenylmethyl chloroformate (Fmoc-Cl) or 9-fluorenylmethyl-N-succinimidyl carbonate (Fmoc-OSu), preferably Fmoc-OSu to furnish 109. The reaction is conveniently run at a temperature of between 0 °C and room temperature, preferably at about room temperature. The compound of formula 109 is a species of amino acid of Structure 5 in Reaction Scheme A.

Pharmaceutical Compositions

[0110] The compounds of formulas I and II as well as Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Ala-Trp-Lys-NH₂; and Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2S,3S) β -methyl-Trp-Lys-NH₂ prepared in accordance with this invention may be prepared in pharmaceutical compositions suitable for administration by injection with a suitable carrier or vehicle by methods known in the art.

Treating Obesity

[0111] The compounds prepared in accordance with this invention exhibited selective MC-4 receptor agonist activity *in vitro*. It is known that agonists of MC4-R activity cause


[illegible]

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523</
--	---	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-------

EXAMPLE 1


Preparation of Fmoc-1-amino-4-phenylcyclohexane-1-carboxylic acid (Fmoc-Apc)

Step 1:



[0113] To a solution of 4-phenylcyclohexanone (10.0 g, 57.5 mmol) in ethanol (100 mL) and water (33 mL) in a glass pressure bottle, were added ammonium carbonate (33 g, 344 mmol, 6 equiv.) and potassium cyanide (5.6 g, 86.2 mmol, 1.5 equiv.). The mixture was heated at 80-90 °C for 24 hrs. The cooled reaction mixture was added to icy water (400 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin A as a white solid (14.0 g, 100% yield). ¹H NMR (DMSO-d₆): 8.63 (s, 1H), 7.23-7.36 (m, 4), 7.15 (m, 1), 2.50 (m, 1H), 2.10 (m, 1H), 1.85 (d, 1H) and 1.55-1.80 (m, 6H).


Step 2:



[0114] The hydantoin A (10.0 g) was suspended in aqueous NaOH (6N, 350 mL) and heated at 130 °C for 2-3 days. Upon the completion of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH ~6). The resulting slurry was filtered, washed with water and dried to give 1-amino-4-phenylcyclohexane carboxylic acid (APC) as a white solid (25 g, >100 % yield. contaminated with inorganic salt) which was

used directly for next step. Small portion of the crude product was purified on HPLC. ^1H NMR (DMSO-d_6): 7.23~7.735 (m, 2), 7.10-7.19 (m, 3H), 2.45 (m, 1H), 1.92-2.18 (m, 3H), 1.56-1.78 (m, 4H) and 1.20 (m, 1H); LRMS (electrospray) m/e 220 ($\text{M}+1$)⁺, Calcd for $\text{C}_{13}\text{H}_{17}\text{NO}_2$, 219.

Step 3:




[0115] The crude 1-amino-4-phenylcyclohexane carboxylic acid (APC) from the last step (25 g) was suspended in dioxane (300 mL) and aqueous 10 % Na_2CO_3 (150 ml) and stirred vigorously overnight. The reaction mixture was concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 5-6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na_2SO_4 . Removal of the solvent gave the crude product which was then purified on flash chromatography (hexane/EtOAc to $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to give pure Fmoc-*cis*-APC (18.2 g, 72% overall yield for two steps) and Fmoc-*trans*-APC (2.1 g, 8 %). Fmoc-*cis*-APC, ^1H NMR(CD_3OD), 7.79 (d, 2H), 7.72 (d, 2H), 7.37 (t, 2), 7.24-7.32 (m, 4), 7.14-7.23 (m, 3), 4.37 (d, 2H), 4.24 (t, 1H), 2.55 (m, 1H), 2.28 (m, 2H), 1.84-1.96 (m, 2H) and 1.64-1.73 (m, 4H).

EXAMPLE 2

Preparation of Fmoc-1-amino-4-(4-methoxyphenyl)cyclohexane-1-carboxylic acid (Fmoc-4-MeOApC-OH)


Step 1:



[0116] A solution of 4-(4-hydroxyphenyl)cyclohexanone (5.0 g, 26.3 mmol) in acetone (100 mL) was treated with K_2CO_3 (14.5 g, 105 mmol, 4 equiv) and iodomethane (4.9 mL, 11.2


g, 78.6 mmol, 3 equiv.). The reaction was heated at 65 °C overnight. After the solvent was removed, the residue was treated with H₂O and extracted with EtOAc. The organic extracts were combined and washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give the spectroscopically pure 4-(4-methoxyphenyl)-cyclohexanone (5.34 g, 100 %). ¹H NMR(CDCl₃) 7.16 (dt, 2H), 6.87 (dt, 2H), 3.78 (s, 3H), 2.99 (tt, 1H), 2.47-2.53 (m, 4H), 2.20 (m, 2H) and 1.83-1.98 (m, 2H); MS (electrospray) *m/e*, 205 (M+1)⁺, Calcd for C₁₃H₁₆O₂, 204.

Step 2:



[0117] To a solution of the 4-(4-methoxyphenyl)-cyclohexanone (3.86 g, 18.9 mmol) in ethanol (50 mL) and water (15 mL) in a glass pressure bottle, were added ammonium carbonate (14.5 g, 151 mmol, 8 equiv.) and potassium cyanide (2.0 g, 30.7 mmol, 1.6 equiv.). The mixture was heated at 80-90 °C for 24 hrs. The cooled reaction mixture was added to icy water (300 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **B** as a white solid (4.75 g, 91% yield). MS (electrospray) *m/e* 273 (M-H), Calcd for C₁₅H₁₈N₂O₃, 274.

Step 3:



[0118] To a suspension of the hydantoin **B** (18.7 g, 68.25 mmol) in dry THF (450 mL) were added di-*tert*-butyl dicarbonate (37.2 g, 170.5 mmol, 2.5 equiv), triethylamine (10.5 mL,

7.59 g, 75.0 mmol, 1.1 equiv) and DMAP (460 mg, 3.65 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (800 mL), washed with 1N HCl (3x50 mL), saturated aqueous Na₂CO₃ (2x50 mL) and brine (2x50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→70/30) to give the pure bis-Boc hydantoin **C** as a white solid (27.6 g, 87%). ¹H NMR (CDCl₃): 7.28 (dt, 2H), 6.88 (dt, 2H), 3.79 (s, 3H), 2.14-2.24 (m, 2H), 1.59 (s, 9H) and 1.38 (s, 9H); MS (electrospray) *m/e* 538 (M+MeCN+Na)⁺, Calcd for C₂₅H₃₄N₂O₇, 474.

Step 4:




[0119] The bis-Boc hydantoin **C** (15.08 g, 31.78 mmol) was dissolved in DME (500 mL) to give a clear solution. To this solution was added 1N NaOH (290 mL, 290 mmol) and the reaction was stirred overnight at room temperature, giving a slightly cloudy mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 1-amino-4-(4-methoxyphenyl)cyclohexane carboxylic acid (4-MeOAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~300 mL) were added DME (300 mL) and a solution of Fmoc-OSu (16.7 g, 49.42 mmol) in DME (200 mL) and the reaction was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove DME, acidified with 3N HCl, extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (CH₂Cl₂/MeOH, 98/2→90/10) to give the pure product

Fmoc-4-MeOAPC as a white solid (12.4 g, 83 % yield from the bis-Boc hydantoin C).
¹H NMR (DMSO-d₆), 7.88 (d, 2H), 7.76 (d, 2H), 7.40 (t, 2H), 7.30 (t, 2H), 7.11 (d, 2H), 6.85 (d, 2H), 3.71 (s, 3H); MS (electrospray) *m/e* 470 (M-H), Calcd for C₂₉H₂₉NO₅, 471.

EXAMPLE 3


Preparation of Fmoc-1-amino-4-(4-ethoxyphenyl)cyclohexane-1-carboxylic acid (**Fmoc-4-EtOapc-OH**)

Step 1:



[0120] A solution of 4-(4-hydroxyphenyl)cyclohexanone (5.0 g, 26.3 mmol) in acetone (100 mL) was treated with K₂CO₃ (14.5 g, 105 mmol, 4 equiv) and iodoethane (10.5 mL, 20.5 g, 131 mmol, 5 equiv.). The reaction was heated at 65 °C overnight. After the solvent was removed, the residue was treated with H₂O and extracted with EtOAc. The organic extracts were combined and washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give the spectroscopically pure 4-(4-ethoxyphenyl)-cyclohexanone (5.74 g, 100 %). ¹H NMR (CDCl₃) 7.15 (dt, 2H), 6.86 (dt, 2H), 4.02 (q, 2H), 2.99 (tt, 1H), 2.46-2.54 (m, 4H), 2.16-2.24 (m, 2H), 1.83-2.00 (m, 2H) and 1.41 (t, 3H); MS (electrospray) *m/e*, 219 (M+1)⁺, Calcd for C₁₄H₁₈O₂, 218.


Step 2:



[0121] To a solution of the 4-(4-ethoxyphenyl)-cyclohexanone (4.15 g, 19.01 mmol) in ethanol (50 mL) and water (15 mL) in a glass pressure bottle, were added ammonium


carbonate (14.5 g, 151 mmol, 8 equiv.) and potassium cyanide (2.05 g, 31.42 mmol, 1.6 equiv.). The mixture was heated at 80-90 °C for 19 hrs. The cooled reaction mixture was added to icy water (300 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **D** as a white solid (5.17 g, 94% yield). MS (electrospray) *m/e* 287 (M-H), Calcd for C₁₆H₂₀N₂O₃, 288.

Step 3:



[0122] To a suspension of the hydantoin **D** (4.22 g, 14.65 mmol) in dry THF (100 mL) were added di-*tert*-butyl dicarbonate (7.98 g, 36.60 mmol, 2.5 equiv), triethylamine (2.3 mL, 1.63 g, 16.11 mmol, 1.1 equiv) and DMAP (89.4 mg, 0.73 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (300 mL), washed with 1N HCl (3x20 mL), saturated aqueous Na₂CO₃ (2x20 mL) and brine (2x20 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→70/30) to give the pure bis-Boc hydantoin **E** as a white solid (7.01 g, 98%). ¹H NMR (CDCl₃): 7.27 (dt, 2H), 6.87 (dt, 2H), 4.02 (q, 2H), 1.59 (s, 9H), 1.43 (t, 3H) and 1.38 (s, 9H); MS (electrospray) *m/e* 999 (2M+Na)⁺, Calcd for C₂₆H₃₆N₂O₇, 488.

Step 4:




[0123] The bis-Boc hydantoin **E** (6.58 g, 13.46 mmol) was dissolved in DME (200 mL) to give a clear solution. To this solution was added 1N NaOH (121 mL, 121 mmol) and the reaction was stirred overnight at room temperature, giving a slightly cloudy mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 1-amino-4-(4-ethoxyphenyl)cyclohexane carboxylic acid (4-EtOAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~130 mL) were added DME (100 mL) and a solution of Fmoc-OSu (6.83 g, 20.24 mmol) in DME (30 mL) and the reaction was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove DME, acidified with 3N HCl, extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (CH₂Cl₂/MeOH, 98/2→90/10) to give the pure product Fmoc-4-EtOAPC as a white solid (5.56g, 85 % yield from the bis-Boc hydantoin **E**). ¹H NMR (DMSO-d₆), 7.88 (d, 2H), 7.74 (d, 2H), 7.40 (td, 2H), 7.30 (td, 2H), 7.11 (d, 2H), 6.84 (d, 2H), 3.97 (q, 2H) and 1.29 (t, 3H); MS (electrospray) *m/e* 484 (M-H), Calcd for C₃₀H₃₁NO₅, 485.

EXAMPLE 4


Preparation of Fmoc-1-amino-4-(4-hydroxyphenyl)cyclohexane-1-carboxylic acid (**Fmoc-4-HOAPc-OH**)

Step 1:




[0124] To a solution of 4-(4-hydroxyphenyl)cyclohexanone (2.00 g, 10.52 mmol) in ethanol (30 mL) and water (10 mL) in a glass pressure bottle, were added ammonium carbonate (6.17 g, 64.2 mmol, 6 equiv.) and potassium cyanide (1.07 g, 15.8 mmol, 1.5 equiv.). The mixture was heated at 80-90 °C overnight. The cooled reaction mixture was added to icy water (200 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **F** as a white solid (2.56 g, 94% yield). MS (electrospray) m/e 261 (M+H)⁺, Calcd for C₁₄H₁₆N₂O₃, 260.

Step 2:



[0125] The hydantoin **F** (2.10 g, 8.06 mmol) was suspended in aqueous NaOH (6N, 100 mL) and heated at 130 °C for 2-3 days. Upon the completion of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH~6). The resulting slurry was filtered, washed with water and dried to give 1-amino-4-(4-hydroxyphenyl)cyclohexane carboxylic acid (4-HOAPC) as a white solid (3.1 g, >100 % yield, contaminated with inorganic salt). MS (electrospray) m/e 236 (M+H)⁺, Calcd for C₁₃H₁₇NO₃, 235.

Step 3:




[0126] The crude 1-amino-4-(4-hydroxyphenyl)cyclohexane carboxylic acid (4-HOAPC) from the last step (3.1 g) was suspended in dioxane (100 mL) and aqueous 10 % Na₂CO₃ (50 ml) and stirred vigorously overnight. The reaction mixture was concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 5-6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na₂SO₄. Removal of the solvent gave the crude product which was purified on flash chromatography (hexane/EtOAc to CH₂Cl₂/MeOH) to give pure Fmoc-4-HOAPC (2.76 g, 75% overall yield for two steps). ¹H NMR(CD₃OD), 7.78 (d, 2H), 7.72 (d, 2H), 7.38 (t, 2H), 7.30 (td, 2H), 7.04 (d, 2H), 6.72 (dt, 2H), 4.38 (d, 2H), 4.25 (t, 1H), 2.46 (m, 1H), 2.24-2.34 (m, 2H) and 1.81-1.92 (m, 6H); MS (electrospray) *m/e* 456 (M-H), Calcd for C₂₈H₂₇NO₅, 457.

EXAMPLE 5

Preparation of Fmoc-1-amino-4-(4-isopropoxyphenyl)cyclohexane-1-carboxylic acid (Fmoc-4-iPrOApc-OH)


Step 1:



[0127] A solution of 4-(4-hydroxyphenyl)cyclohexanone (6.0 g, 31.6 mmol) in DMF (90 mL) was treated with K₂CO₃ (21 g, 158 mmol, 5 equiv) and 2-iodopropane (15 mL, 26.8 g, 158 mmol, 5 equiv.). The reaction was heated at 100 °C overnight. After the solvent was removed, the residue was treated with H₂O and extracted with EtOAc. The organic


extracts were combined and washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give the spectroscopically pure 4-(4-isopropoxyphenyl)cyclohexanone (7.02 g, 95 %). ¹H NMR (CDCl₃): 7.14 (dt, 2H), 6.84 (dt, 2H), 4.3 (septet, 1H), 2.97 (tt, 1H), 2.46-2.52 (m, 4H), 2.16-2.24 (m, 2H), 1.83-1.98 (m, 2H) and 1.33 (d, 6H).

Step 2:



[0128] To a solution of the 4-(4-isopropoxyphenyl)cyclohexanone (5.1 g, 21.98 mmol) in ethanol (90 mL) and water (30 mL) in a glass pressure bottle, were added ammonium carbonate (12.6 g, 131 mmol, 6 equiv.) and potassium cyanide (2.14 g, 32.9 mmol, 1.5 equiv.). The mixture was heated at 80-90 °C for 24 hrs. The cooled reaction mixture was added to icy water (400 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield hydantoin **G** as a white solid (6.60 g, 99% yield). ¹H NMR (DMSO-d₆): 10.60 (s, 1H), 8.65 (s, 1H), 7.18 (d, 2H), 6.80 (d, 2H), 4.52 (septet, 1H), 2.43 (m, 1H), 1.85-2.15 (m, 2H), 1.56-1.80 (m, 6H) and 1.22 (d, 6H); MS (electrospray) *m/e* 301 (M-H), Calcd for C₁₇H₂₂N₂O₃, 302.


Step 3:



[0129] To a suspension of the hydantoin **G** (5.8 g, 19.20 mmol) in dry THF (180 mL) were added di-*tert*-butyl dicarbonate (10.46 g, 48.0 mmol, 2.5 equiv), triethylamine (2.9 mL, 2.13 g, 21.12 mmol, 1.1 equiv) and DMAP (140 mg, 1.15 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was

stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (600 mL), washed with 1N HCl (3x40 mL), saturated aqueous Na₂CO₃ (2x40 mL) and brine (2x40 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→80/20) to give the pure bis-Boc hydantoin **H** as a white solid (9.4 g, 98%). ¹H NMR (CDCl₃): 7.27 (dt, 2H), 6.87 (dt, 2H), 4.02 (q, 2H), 2.98 (t, 1H), 2.26-2.56 (m, 4H), 2.14-2.24 (m, 2H), 1.76-1.86 (m, 2H), 1.59 (s, 9H), 1.43 (t, 3H) and 1.38 (s, 9H); MS (electrospray) *m/e* 999 (2M+Na)⁺, Calcd for C₂₆H₃₆N₂O₇, 488.

Step 4:




[0130] The bis-Boc hydantoin **H** (4.34 g, 8.64 mmol) was dissolved in DME (100 mL) to give a clear solution. To this solution was added 1N NaOH (78 mL, 78 mmol) and the reaction was stirred overnight at room temperature, giving a fairly clear mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 1-amino-4-(4-isopropoxyphenyl)cyclohexane carboxylic acid (4-iPrOAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~90 mL) were added DME (120 mL) and a solution of Fmoc-OSu (3.49 g, 10.34 mmol, 1.2 equiv) in DME (20 mL) and the reaction was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove DME, acidified with 3N HCl, extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (hexane/EtOAc→CH₂Cl₂/MeOH) to give the pure product Fmoc-4-iPrOAPC as a white solid (3.23, 75 % yield from bis-Boc

hydantoin **H**). ^1H NMR(DMSO- d_6), 7.76 (d, 2H), 7.60 (d, 2H), 7.39 (t, 2H), 7.31 (t, 2H), 7.08 (d, 2H), 6.84 (d, 2H), 4.24 (m, 1H) and 1.34 (d, 6H); MS (electrospray) m/e 498 (M-H), Calcd for $\text{C}_{31}\text{H}_{33}\text{NO}_5$, 499.

EXAMPLE 6


Preparation of Fmoc-1-amino-4-(4-methylphenyl)cyclohexane-1-carboxylic acid (**Fmoc-4-MeApc-OH**)

Step 1:




[0131] To a solution of 4-iodotoluene (10.9 g, 50.0 mmol) in dry THF (180 mL) at -78°C was added a solution of $n\text{-BuLi}$ (1.6 M, 31.0 mL, 50 mmol) in hexane over 20 min. The reaction was stirred for another 20 min before a solution of 1,4-cyclohexanedione *mono*-ethylene ketal (6.0 g, 38.46 mmol) in dry THF (100 mL) was added dropwise. After stirred for 2 h at -78°C , the reaction was quenched with aqueous NH_4Cl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na_2SO_4 , concentrated in vacuo to give the spectroscopically pure product **I** as a white solid (9.34 g, 98% yield). ^1H NMR (CDCl_3): 7.41 (m, 2H), 7.16 (d, 2H), 3.98 (m, 4H), 2.34 (s, 3H); MS (EI) m/e 248 (M^+), Calcd for $\text{C}_{15}\text{H}_{20}\text{O}_3$, 248.

Step 2:




[0132] To a solution of the alcohol **I** (9.10g, 36.65 mmol) in dry benzene (200 mL) in a flask equipped with a Dean-Stark trap, was added p-toluenesulfonic acid monohydrate (650 mg) and the reaction was heated at 100 °C for 3 hrs. The reaction was cooled to rt, diluted with EtOAc (500 mL) and washed with aqueous Na₂CO₃ (50 mL), brine (3x50 mL), dried over Na₂SO₄ and concentrated under reduced pressure to give the spectroscopically pure product **J** (8.36 g, 100 yield), which was used for next step without purification. MS (EI) *m/e* 230 (M⁺), 190 (M-OCH₂CH₂O), Calcd for C₁₅H₁₈O₂, 230.

Step 3:



[0133] To a solution of the olefin **J** (7.49 g) in EtOAc (180 mL) was added Pd/C (5 wt % on carbon, 800 mg) and the reaction was run under 40 psi of hydrogen for 3 hrs at room temperature. The catalyst was filtered off and the filtrate was concentrated to give the spectroscopically pure product **K** as a colorless oil (7.40 g, 100% yield). MS (EI) *m/e* 232 (M⁺), 188 (M-OCH₂CH₂), Calcd for C₁₅H₂₀O₂, 232.


Step 4:



[0134] A solution of the ketal **K** (6.90 g) in acetone (140 mL) was treated with 4N HCl (60 mL) and heated at 65 °C for 4 hrs. Solvent was removed and the residue was diluted with EtOAc and neutralized with 4N HCl. The aqueous was extracted with EtOAc. The


combined organic extracts were washed with brine, dried and concentrated. The resulting crude 4-(4-methylphenyl)cyclohexanone was used for next step without purification (5.57 g, quantitative yield). MS (EI) m/e 188 (M^+), Calcd for $C_{13}H_{16}O$, 188.

Step 5:



[0135] To a solution of 4-(4-methylphenyl)cyclohexanone (5.32 g, 28.3 mmol) in ethanol (90 mL) and water (30 mL) in a glass pressure bottle, were added ammonium carbonate (16.3 g, 169.8 mmol, 6 equiv.) and potassium cyanide (3.68 g, 56.5 mmol, 2 equiv.). The mixture was heated at 80-90 °C overnight. The cooled reaction mixture was added to icy water (400 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **L** as a white solid (6.3 g, 86% yield). MS (electrospray) m/e 517 (2M+H), Calcd for $C_{15}H_{18}ClN_2O_2$, 258


Step 6:



[0136] To a suspension of the hydantoin **L** (5.82 g, 22.55 mmol) in dry THF (250 mL) were added di-*tert*-butyl dicarbonate (12.3 g, 56.4 mmol, 2.5 equiv), triethylamine (3.5 mL, 2.5 g, 24.7 mmol, 1.1 equiv) and DMAP (275 mg, 2.25 mmol) in succession. The reaction turned into a clear yellow solution and was stirred overnight at room temperature. The

reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (500 mL), washed with 1N HCl (3x50 mL), saturated aqueous Na₂CO₃ (2x50 mL) and brine (2x50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→70/30) to give the pure bis-Boc hydantoin **M** as a white solid (10.03 g, 100% yield). ¹H NMR (CDCl₃): 7.26 (d, 2H), 6.87 (d, 2H), 3.00 (m, 1H), 2.32 (s, 3H), 1.59 (s, 9H) and 1.37 (s, 9H).

Step 7:




[0137] The bis-Boc hydantoin **M** (6.40 g, 13.97 mmol) was dissolved in DME (200 mL) to give a clear solution. To this solution was added 1N NaOH (120 mL, 120 mmol) and the reaction was stirred overnight at room temperature, giving a slightly cloudy mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 1-amino-4-(4-methylphenyl)cyclohexane carboxylic acid (4-MeAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~140 mL) were added DME (240 mL) and a solution of Fmoc-OSu (5.10 g, 15.13 mmol, 1.1 equiv) in DME (40 mL) and the reaction was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove DME, acidified with 3N HCl, extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (CH₂Cl₂/MeOH, 98/2→90/10) to give the pure product Fmoc-4-MeAPC as a white solid (4.35 g, 69 % yield from bis-Boc hydantoin **M**). ¹H NMR (DMSO-d₆): 7.88 (d, 2H), 7.75 (d, 2H), 7.24-7.43 (m, 4H), 7.02-7.14 (m, 4H), 4.25 (m, 3H), 2.24 (s, 3H).

EXAMPLE 7


Preparation of Fmoc-1-amino-4-(4-chlorophenyl)cyclohexane-1-carboxylic acid (**Fmoc-4-ClApc-OH**)

Step 1:




[0138] A solution of 4-chlorophenylbromide (7.5 g, 39.2 mmol) in dry THF (180 mL) was cooled to -78°C and treated dropwise with a solution of n-BuLi (1.6 M, 25 mL, 40 mmol) in hexane over 20 min. The reaction was stirred for a further 30 min before a solution of 1,4-cyclohexanedione *mono*-ethylene ketal (6.0 g, 38.46 mmol) in dry THF (100 mL) was added dropwise. After stirred for 1 hr at -78°C , the reaction was quenched with aqueous NH_4Cl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na_2SO_4 , concentrated in vacuo to give the spectroscopically pure product **N** as a white solid (9.40 g, 91% yield). ^1H NMR (CDCl_3): 7.45 (m, 2H), 7.31 (m, 2H), 3.99 (m, 4H), 2.02-2.20 (m, 4H), 1.75-1.82 (m, 2H), 1.66-1.73 (m, 2H), 1.54 (s, 1H); MS (EI) m/e 268 (M^+), 251 (M-OH), 250 (M- H_2O), Calcd for $\text{C}_{14}\text{H}_{17}\text{ClO}_3$, 268.

Step 2:




[0139] To a solution of the alcohol **N** (6.78 g, 25.30 mmol) in dry benzene (120 mL) in a flask equipped with a Dean-Stark trap, was added p-toluenesulfonic acid monohydrate (960 mg) and the reaction was heated at reflux for 3 hrs. The reaction was cooled to rt, diluted with EtOAc (500 mL) and washed with aqueous Na₂CO₃ (50 mL), brine (3x50 mL), dried over Na₂SO₄ and concentrated under reduced pressure to give the spectroscopically pure product **O** (6.30 g, 100 yield), which was used for next step without purification. MS (EI) *m/e* 250 (M⁺), 190 (M-OCH₂CH₂O), Calcd for C₁₄H₁₅ClO₂, 250.

Step 3:



[0140] To a solution of the olefin **O** (6.11 g) in EtOAc (120 mL) was added Pd/C (5 wt % on carbon, 600 mg) and the reaction was run under 5 psi of hydrogen for 3 hrs at room temperature. The catalyst was filtered off and the filtrate was concentrated to give the spectroscopically pure product **P** as a colorless oil (6.10 g, 100% yield). MS (EI) *m/e* 252(M⁺), Calcd for C₁₄H₁₇ClO₂, 252.


Step 4:



[0141] A solution of the ketal **P** (5.81 g, 23.06 mmol) in acetone (200 mL) was treated with p-toluenesulfonic acid monohydrate (876 mg) and heated at 60 °C overnight. Solvent was removed and the residue was taken up in EtOAc, washed with aqueous Na₂CO₃ solution, brine, dried and concentrated to give the crude product as a yellow oil (5.38 g, >100%


yield). Purification through flash chromatography (hexane/EtOAc, 80/20→60/40) provided 4-(4-chlorophenyl)cyclohexanone as a light yellow oil (4.54 g, 95% yield). MS (EI) m/e 208 (M^+), Calcd for $C_{12}H_{13}ClO_2$, 208.

Step 5:



[0142] To a solution of 4-(4-chlorophenyl)cyclohexanone (4.26 g, 20.48 mmol) in ethanol (90 mL) and water (30 mL) in a glass pressure bottle, were added ammonium carbonate (13.8 g, 144 mmol, 7 equiv) and potassium cyanide (3.56 g, 54.77 mmol, 2.5 equiv). The mixture was heated at 80-90 °C overnight. The cooled reaction mixture was added to icy water (400 mL) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **Q** as a white solid (5.58 g, 98% yield). MS (electrospray) m/e 277 ($M-H$), Calcd for $C_{14}H_{15}ClN_2O_2$, 278.


Step 6:



[0143] To a suspension of the hydantoin **Q** (5.15 g, 18.5 mmol) in dry THF (250 mL) were added di-*tert*-butyl dicarbonate (10.1 g, 46.3 mmol, 2.5 equiv), triethylamine (2.8 mL, 2.07 g, 20.45 mmol, 1.1 equiv) and DMAP (226 mg, 1.85 mmol) in succession. The reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (500 mL), washed with 1N HCl (3x50 mL),

saturated aqueous Na_2CO_3 (2x50 mL) and brine (2x50 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→70/30) to give the pure bis-Boc hydantoin **R** as a white solid (8.05 g, 91% yield). MS (electrospray) m/e 542 ($\text{M}+\text{Ma}+\text{MeCN}$), Calcd for $\text{C}_{24}\text{H}_{31}\text{ClN}_2\text{O}_6$, 478.

Step 7:




[0144] The bis-Boc hydantoin **R** (6.41 g, 13.97 mmol) was dissolved in DME (200 mL) to give a clear solution. To this solution was added 1N NaOH (120 mL, 120 mmol) and the reaction was stirred overnight at room temperature, giving a slightly cloudy mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et_2O . Without purification, the resulting aqueous layer containing 1-amino-4-(4-chlorophenyl)cyclohexane carboxylic acid (4-ClAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~180 mL) were added DME (240 mL) and a solution of Fmoc-OSu (5.31 g, 15.74 mmol, 1.1 equiv) in DME (30 mL) and the reaction was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove DME, acidified with 3N HCl, extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The crude product was purified through flash chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 98/2→90/10) to give the pure product Fmoc-4-ClAPC as a white solid (5.04 g, 76% yield from the bis-Boc hydantoin). ^1H NMR ($\text{DMSO}-d_6$), 7.88 (d, 2H), 7.74 (d, 2H), 7.19-7.42 (m, 8H), 4.20-4.31 (m, 3H); MS (electrospray) m/e 474 ($\text{M}-\text{H}$), Calcd for $\text{C}_{28}\text{H}_{26}\text{ClNO}_4$, 475.

EXAMPLE 8


Preparation of Fmoc-1-amino-4-(3-methoxyphenyl)cyclohexane-1-carboxylic acid (**Fmoc-3-MeOApc-OH**)

Step 1:



[0145] To a solution of 3-iodoanisole (11.7, 50.0 mmol, 1.3 equiv) in dry THF (180 mL) at -78 °C was added a solution of n-BuLi (1.6 M, 31.0 mL, 50 mmol, 1.3 equiv) in hexane over 25 min. The reaction was stirred for another 30 min before a solution of 1,4-cyclohexanedione *mono*-ethylene ketal (6.0 g, 38.46 mmol) in dry THF (100 mL) was added dropwise. After stirred for 2 h at -78 °C, the reaction was quenched with aqueous NH₄Cl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄, concentrated in vacuo to give the spectroscopically pure product **S** as a white solid (9.34 g, 98% yield). ¹H NMR (CDCl₃): 7.26 (dd, 1H), 7.06-7.11 (m, 2H), 6.79 (dd, 1H), 3.98 (m, 4H), 3.81 (s, 3H).


Step 2:



[0146] To a stirred solution of the alcohol **S** (5.6 g, 21.21 mmol) in dry CH₂Cl₂ (200 mL) under a nitrogen atmosphere at salt-ice bath temperature, were added in succession triethylsilane (10.2 mL, 7.4 g, 63.63 mmol, 3 equiv) and boron trifluoride etherate (21.5


mL, 24.1 g, 169.7 mmol, 8 equiv). The reaction mixture was then allowed to warm to room temperature and stirred for 3 hrs before washed with 10% aqueous K_2CO_3 solution and H_2O , dried over Na_2SO_4 , and concentrated in vacuo to give the deoxygenation compound as an oil (4.91g), which was sufficiently pure for direct use. This crude intermediate was dissolved in acetone (130 mL) and treated with 4N HCl (60 mL) and heated at 65 °C for 4 hrs. Solvent was removed under reduced pressure and the residue was diluted with EtOAc and neutralized with 4N NaOH solution. The aqueous layer was extracted with EtOAc and the combined organic extracts were washed with brine, dried and concentrated. The resulting residue was purified by flash chromatography on silica gel (80/20→60/40) to give 4-(3-methoxyphenyl)cyclohexanone (3.67 g, 85% overall yield) as a yellow oil. 1H NMR ($CDCl_3$): 7.25 (dt, 1H), 6.75-6.86 (m, 3H), 3.81 (s, 3H), 3.00 (tt, 1H); MS (EI) m/e 204 (M^+), Calcd for $C_{13}H_{16}O_2$, 204.

Step 3:




[0147] To a solution of 4-(3-methoxyphenyl)cyclohexanone (3.10 g, 15.20 mmol) in ethanol (60 mL) and water (20 mL) in a glass pressure bottle, were added ammonium carbonate (8.75 g, 91.20 mmol, 6 equiv.) and potassium cyanide (1.98 g, 30.40 mmol, 2 equiv.). The mixture was heated at 80-90 °C overnight. The cooled reaction mixture was added to icy water (300 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin T as a white solid (4.08 g, 98% yield). 1H NMR ($DMSO-d_6$): 7.11 (d, 1H), 6.70-6.94 (m, 3H), 3.72 (s, 3H); MS (electrospray) m/e 316 ($M+MeCN+H$), Calcd for $C_{15}H_{18}N_2O_3$, 274.

Step 4:



[0148] To a suspension of the hydantoin **T** (5.29 g, 19.30 mmol) in dry THF (250 mL) were added di-*tert*-butyl dicarbonate (10.5 g, 48.16 mmol, 2.5 equiv), triethylamine (3.0 mL, 2.17 g, 21.52 mmol, 1.1 equiv) and DMAP (235 mg, 1.92 mmol) in succession. The reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (500 mL), washed with 1N HCl (3x50 mL), saturated aqueous Na₂CO₃ (2x50 mL) and brine (2x50 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 80/20→60/40) to give the pure bis-Boc hydantoin **U** as a white solid (8.70 g, 95% yield). MS (electrospray) *m/e* 538 (M+MeCN+Na), Calcd for C₂₅H₃₄N₂O₇, 474.

Step 5:




[0149] The bis-Boc hydantoin **U** (2.30 g, 4.84 mmol) was dissolved in DME (80 mL) to give a clear solution. To this solution was added 1N NaOH (44 mL, 44 mmol) and the reaction was stirred overnight at room temperature, giving a slightly cloudy mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under

reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 1-amino-4-(3-methoxyphenyl)cyclohexane carboxylic acid (3-MeOAPC) was treated with 6N HCl to adjust the pH to 11-12. To this solution (~40 mL) were added dioxane (80 mL) and Fmoc-Cl (1.73 g, 6.76 mmol, 1.4 equiv) and the reaction was stirred overnight at room temperature. The reaction mixture was then concentrated under reduced pressure to remove DME, neutralized with 3N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified by flash chromatography on silica gel (CH₂Cl₂/MeOH, 98/2→90/10) to give Fmoc-3-MeOAPC as a white solid (1.98 g, 87 % yield from bis-Boc hydantoin U). ¹H NMR (DMSO-d₆), 7.88 (d, 2H), 7.75 (d, 2H), 7.40 (td, 2H), 7.30 (td, 2H), 7.21 (m, 1H), 6.71-6.80 (m, 3H), 3.72 (s, 3H); MS (electrospray) *m/e* 494 (M+Na), Calcd for C₂₉H₂₉NO₅, 471.

EXAMPLE 28

Preparation of Fmoc-(D,L)-5-bromo-2 aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-Br-Atc-OH**)

Step 1:




[0150] A mixture of 3-(2-bromophenyl)propanoic acid (prepared in 2 steps from 2-bromobenzyl bromide, 2.0 g, 8.73 mmole), oxalyl chloride (1.14 ml, 13.1 mmole) and methylene chloride (20 ml) was cooled in an ice bath and N,N-dimethylformamide (34 μL, 0.44 mmole) was added dropwise. The mixture was stirred at room temperature for 3 hours. Concentration in vacuo gave 3-(2-bromophenyl)propanoyl chloride which was taken up in methylene chloride and used in the next step as a crude.

Step 2:



[0151] A solution of the above acid chloride (crude, 8.73 mmole) in methylene chloride was slowly added to a solution of diazomethane (generated from 5.70 g of 1-methyl-3-nitro-1-nitrosoguanidine) in ether (40 ml) cooled in an ice bath. The mixture was then warmed up to room temperature and stirred overnight. The mixture was concentrated in vacuo and purified by column chromatography (10 \rightarrow 20% ethyl acetate/hexanes) to give 1-diazo-4-(2-bromophenyl)butan-2-one (1.88 g, 85% over 2 steps). ^1H NMR (CDCl_3) δ 7.50 (1H, d, phenyl), 7.24 (2H, m, phenyl), 7.06 (1H, m, phenyl), 5.21 (1H, broad s, diazo), 3.05 (2H, t, benzylic), 2.62 (2H, m).


Step 3:



[0152] To a mixture of rhodium (II) acetate dimer (15 mg, 0.068 mmole) in methylene chloride (120 ml) under reflux was slowly added a solution of 1-diazo-4-(2-bromophenyl)butan-2-one (1.74 g, 6.85 mmole) in methylene chloride (30 ml). After the addition was complete, the mixture was refluxed for an extra twenty minutes. The mixture was cooled to room temperature, trifluoroacetic acid (1.5 ml) was added and the mixture was stirred at room temperature for an hour. The reaction was quenched with saturated sodium bicarbonate solution. The layers were separated and the methylene chloride layer was washed once more with saturated sodium bicarbonate solution. The combined aqueous layers were back-extracted with methylene chloride. The combined


organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo to give a brown oil. Purification by column chromatography (10 → 15% ethyl acetate/hexanes) gave 5-bromo-β-tetralone (1.18 g, 77% yield) as a colorless oil. ¹H NMR (CDCl₃) δ 7.46 (1H, t, phenyl), 7.05-7.09 (2H, m, phenyl), 3.58 (2H, s, benzylic), 3.22 (2H, t, benzylic), 2.54 (2H, t).

Step 4:




[0153] A mixture of 5-bromo-β-tetralone (1.18 g, 5.24 mmole), potassium cyanide (512 mg, 7.86 mmole), ammonium carbonate (3.0 g, 31.22 mmole), ethanol (25 ml) and water (5 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 4 days. After cooling to room temperature, the white slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave hydantoin **V** (1.31 g, 85%). ¹H NMR (DMSO-d₆) δ 10.71 (1H, broad, NH), 8.28 (1H, broad s, NH), 7.0-7.5 (3H, m, phenyl). LRMS (Electrospray): C₁₂H₁₁BrN₂O₂, calc. 294; observed: 293 (M-H), 295 (M-H).

Step 5:



[0154] A mixture of hydantoin V (1.287 g, 4.36 mmole), Ba(OH)₂ · H₂O (4.20 g, 22.2 mmole) in water (25 ml) in a sealed, thick walled pressure flask was heated in a 125°C oil bath for 4 days. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for one hour and cooled to room temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 20 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitate which were filtered, washed with water and dried in vacuo overnight to give racemic 5-bromo-2-aminotetraline-2-carboxylic acid (893 mg, 76% yield). LRMS (Electrospray): C₁₁H₁₂BrNO₂, calc. 269; observed: 270 (M+H), 272 (M+H), 268 (M-H), 270 (M-H).

Step 6:




[0155] A mixture of racemic 5-bromo-2 aminotetraline-2-carboxylic acid (882 mg, 3.27 mmole), triethylamine (0.60 ml, 4.30 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 1.32 g, 3.91 mmole) in acetonitrile (30 ml) and water (30 ml) was stirred at room temperature overnight. TLC analysis of the reaction the next day indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (0.25 g), triethylamine (0.6 ml) and acetonitrile (5 ml) was added and the mixture was stirred at room temperature for another day. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted twice with methylene chloride. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 2 → 5 → 10% methanol/methylene chloride) to give racemic Fmoc-5-bromo-2

aminotetraline-2-carboxylic acid (1.09 g, 68% yield) as a white solid. HRMS (FAB): $C_{26}H_{22}BrNNaO_4$ (M+Na) calc. 514.0630; observed: 514.0643.

EXAMPLE 10

Preparation of Fmoc-(D,L)-5-chloro-2 aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-ClAtc-OH**)

Step 1:



[0156] A mixture of 3-(2-chlorophenyl)propanoic acid (5.0 g, 27.1 mmole), thionyl chloride (10.9 ml, 149 mmole) and toluene (75 ml) was refluxed for two hours. Concentration in vacuo gave 3-(2-chlorophenyl)propanoyl chloride which was taken up in methylene chloride and used in the next step without further purification.


Step 2:



[0157] A solution of the above acid chloride (crude, 27.1 mmole) in methylene chloride was slowly added to a solution of diazomethane (generated from 17.8 g of 1-methyl-3-nitro-1-nitrosoguanidine) in ether (120 ml) cooled in an ice bath. The mixture was then warmed up to room temperature and stirred overnight. The mixture was concentrated in vacuo to give 1-diazo-4-(2-chlorophenyl)butan-2-one (5.87 g, > 100% over 2 steps) as a bright yellow oil. The compound was used in the next step without further purification. 1H


NMR (CDCl₃) δ 7.05-7.32 (4H, m, phenyl), 5.13 (1H, broad s, diazo), 3.00 (2H, t, benzylic), 2.57 (2H, m).

Step 3



[0158] To a mixture of rhodium (II) acetate dimer (60 mg, 0.27 mmole) in methylene chloride (400 ml) under reflux was slowly added a solution of crude 1-diazo-4-(2-bromophenyl)butan-2-one (5.87 g, 27.1 mmole theoretical) in methylene chloride (50 ml). After the addition was complete, the mixture was refluxed for an extra twenty minutes. The mixture was cooled to room temperature, trifluoroacetic acid (6.0 ml) was added and the mixture was stirred at room temperature for two hours. The reaction was quenched with saturated sodium bicarbonate solution. The layers were separated and the methylene chloride layer was washed once more with saturated sodium bicarbonate solution. The combined aqueous layers were back-extracted with methylene chloride. The combined organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo to give a brown oil. Purification by column chromatography (10 → 15% ethyl acetate/hexanes) gave 5-chloro-β-tetralone (3.32 g, 68% yield for steps 1 through 3) as a light brown oil. ¹H NMR (CDCl₃) δ 7.30 (1H, m, phenyl), 7.15 (1H, t, phenyl), 7.05 (1H, d, phenyl), 3.60 (2H, s, benzylic), 3.22 (2H, t, benzylic), 2.56 (2H, t).


Step 4:



[0159] A mixture of 5-chloro-β-tetralone (880 mg, 4.87 mmole), potassium cyanide (500 mg, 7.67 mmole), ammonium carbonate (2.85 g, 29.7 mmole), ethanol (24 ml) and water


(6 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 66 hours. After cooling to room temperature, the slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave hydantoin **W** (0.92 g, 75%) as a light beige solid. ¹H NMR (DMSO-d₆) δ 10.70 (1H, broad, NH), 8.25 (1H, broad s, NH), 7.0-7.3 (3H, m, phenyl). LRMS (Electrospray): C₁₂H₁₁ClN₂O₂, calc. 250; observed: 249 (M-H), 251 (M-H).

Step 5:



[0160] A mixture of hydantoin **W** (880 mg, 3.51 mmole), Ba(OH)₂·H₂O (3.40 g, 18.0 mmole) in water (50 ml, too dilute) in a sealed, thick walled pressure flask was heated in a 125°C oil bath for 2 days. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for two hours and cooled to room temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 50 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitate which were filtered, washed with water and dried in vacuo overnight to give racemic 5-chloro-2-aminotetralin-2-carboxylic acid (788 mg, 99% yield). LRMS (Electrospray): C₁₁H₁₂ClNO₂, calc. 225; observed: 226 (M+H), 228 (M+H), 224 (M-H), 226 (M-H).

Step 6:




[0161] A mixture of racemic 5-chloro-2-aminotetraline-2-carboxylic acid (402 mg, 1.78 mmole), triethylamine (0.38 ml, 2.73 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 904 mg, 2.68 mmole) in acetonitrile (20 ml) and water (20 ml) was stirred at room temperature for two days. TLC analysis of the reaction after two days indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (0.12 g) and triethylamine (0.1 ml) was added and the mixture was stirred at room temperature for another day. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted three times with ethyl acetate. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 3 → 6 → 8% methanol/methylene chloride) to give racemic Fmoc-5-chloro-2-aminotetraline-2-carboxylic acid (540 mg, 68% yield) as a white solid. HRMS (EI): C₂₆H₂₂ClNO₄ (M) calc. 447.1237; observed: 447.1234.

EXAMPLE 11

Preparation of Fmoc-(D,L)-5-methoxy-2-aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-MeOAtc-OH**)

Step 1:




[0162] A mixture of racemic 5-methoxy-2-aminotetraline-2-carboxylic acid (prepared according to Obrecht, D. *et. al. Helv. Chim Acta.* **1992**, 75, 1666) (802 mg, 3.62 mmole), triethylamine (0.62 ml, 4.45 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 1.47 g, 4.36 mmole) in acetonitrile (25 ml) and water (25 ml) was stirred at room temperature for 30 hours. TLC analysis of the reaction indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (370 mg) and

triethylamine (0.6 ml) were added and the mixture was stirred at room temperature for another 24 hours. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion was extracted three times with ethyl acetate. The combined organic layers were washed with water, brine and dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 1 → 3 → 5 → 10% methanol/methylene chloride) to give racemic Fmoc-5-methoxy-2-aminotetraline-2-carboxylic acid (1.14 g, 71% yield) as an off-white solid. HRMS (FAB): C₂₇H₂₆NO₅ (M+H) calc. 444.1812; observed: 444.1814.

EXAMPLE 12

Preparation of Fmoc-(D,L)-5-ethoxy-2-aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-EtOAtc-OH**)


Step 1:



[0163] A mixture of 1,6-dihydroxynaphthalene (5.02 g, 31.3 mmole), anhydrous potassium carbonate (52.0 g, 376 mmole), N,N-dimethylformamide (50 ml) and iodoethane (15 ml, 188 mmole) was stirred in a 35°C oil bath for 24 hours. The reaction mixture was filtered and the solid residue was rinsed thoroughly with ethyl ether. The filtrate and the washings were combined and concentrated in vacuo to remove most of the solvents. The brown residue was partitioned between water and ether and the layers were separated. The ether layer was washed with water. The combined aqueous layers were back extracted with ether. The ether extracts were combined, washed with brine and dried over magnesium sulfate. Filtration and concentration gave a crude brown solid (6.74 g, 99% yield). Recrystallization of the crude product from hot methanol gave 1,6-diethoxynaphthalene (4.36 g, 64% yield, first crop) as a light brown solid. ¹H NMR


(CDCl₃) δ 8.20 (1H, d, phenyl), 7.06-7.36 (4H, m, phenyl), 6.66 (1H, dd, phenyl), 4.10-4.23 (4H, 2 sets of q, 2 CH₂), 1.45-1.56 (6H, 2 sets of t, 2 CH₃).

Step 2:




[0164] To a refluxing solution of 1,6-diethoxynaphthalene (4.15 g, 19.2 mmole) in absolute ethanol (100 ml) was carefully added small pieces of sodium metal (6.8 g, 296 mmole) over 60 minutes. The mixture was refluxed for another 90 minutes. TLC indicated the presence of unreacted starting material. Extra sodium metal (1.0 g, 43.5 mmole) was added and the reaction mixture was refluxed for another 60 minutes. The reaction was cooled to room temperature, quenched with water and acidified with concentrated hydrochloric acid. The mixture was concentrated in vacuo to remove most of the ethanol. The aqueous mixture was extracted three times with ether. The combined organic layers were washed with water and dried over sodium sulfate. Filtration and concentration gave a brown solid which was dissolved in 1:1 ethanol/water (200 ml), then *p*-toluenesulfonic acid (400 mg) was added. The mixture was refluxed for 210 minutes. Extra *p*-toluenesulfonic acid (100 mg) was added and the mixture was refluxed for another 60 minutes. After cooling to room temperature, most of the ethanol was removed under reduced pressure. The aqueous mixture was extracted three times with ether and the combined organic layers were washed with water, saturated sodium chloride solution and dried over sodium sulfate. Filtration and concentration gave a brown oil which was purified by column chromatography (7% ethyl acetate/hexanes) to give 5-ethoxy-β-tetralone (2.43 g, 67% yield) as a light yellow oil. ¹H NMR (CDCl₃) δ 7.15 (1H, t, phenyl), 6.76 (1H, d, phenyl), 6.72 (1H, d, phenyl), 4.05 (2H, q, CH₂), 3.56 (2H, s, benzylic), 3.10 (2H, t, benzylic), 2.53 (2H, t), 1.44 (3H, t, CH₃).

Step 3:



[0165] A mixture of 5-ethoxy- β -tetralone (2.23 g, 11.7 mmole), potassium cyanide (1.20 g, 18.4 mmole), ammonium carbonate (6.75 g, 70.2 mmole), ethanol (80 ml) and water (20 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 3 days. After cooling to room temperature, the slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave hydantoin X (2.69 g, 88%) as a beige solid. ^1H NMR (DMSO- d_6) δ 10.65 (1H, broad s, NH), 8.22 (1H, broad s, NH), 7.06 (1H, t, phenyl), 6.75 (1H, d, phenyl), 6.65 (1H, d, phenyl), 3.98 (2H, q, CH_2), 1.32 (3H, t, CH_3). LRMS (Electrospray): $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_3$, calc. 259; observed: 258 (M-H).


Step 4:



[0166] A mixture of hydantoin X (2.57 g, 9.87 mmole), $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ (9.40 g, 49.6 mmole) in water (200 ml, too dilute) in a sealed, thick walled pressure flask was heated in a 105°C oil bath for 39 hours. Extra $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ (9.40 g, 49.6 mmole) was added and the mixture was heated in a 125°C oil bath for an additional 21 hours. The reaction mixture was cooled to room temperature, acidified to $\sim \text{pH}$ 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for one hour and cooled to room temperature. The white suspension was filtered and the precipitates

rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 75 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitate which were filtered, washed with water and air-dried to give racemic 5-ethoxy-2-aminotetraline-2-carboxylic acid (2.34 g, quantitative yield) as a light beige solid. LRMS (Electrospray): $C_{13}H_{17}NO_3$, calc. 235; observed: 236 (M+H), 234 (M-H).

Step 5:



[0167] A mixture of racemic 5-ethoxy-2-aminotetraline-2-carboxylic acid (2.22 g, 9.44 mmole), triethylamine (2.00 ml, 14.3 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 4.81 g, 14.3 mmole) in acetonitrile (75 ml) and water (75 ml) was stirred at room temperature for two days. TLC analysis of the reaction indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (645 mg) and triethylamine (1.0 ml) was added and the mixture was stirred at room temperature for another day. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted three times with ethyl acetate. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 3 → 5 → 10% methanol/methylene chloride) to give racemic Fmoc-5-ethoxy-2-aminotetraline-2-carboxylic acid (4.66 g, > quantitative yield) as a white solid. HRMS (FAB): $C_{28}H_{28}NO_5$ (M+H) calc. 458.1967; observed: 458.1985.

EXAMPLE 13


Preparation of Fmoc-(D,L)-5-isopropoxy-2-aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-iPrOAtc-OH**)

Step 1:



[0168] A mixture of 6-methoxy-1-tetralone (5.07 g, 28.8 mmole), 10% Pd/C (3.53 g, 3.32 mmole) in dry *p*-cymene (250 ml) was heated to reflux under argon for 38 hours. The reaction mixture was cooled to room temperature, filtered over celite and the residue rinsed thoroughly with *p*-cymene. The filtrate and the washings were combined and extracted twice with 1N sodium hydroxide solution (2 x 70 ml). The combined aqueous extracts were acidified with 6N hydrochloric acid to pH ~3 and extracted three times with ether. The combined organic layers were washed with water, dried over anhydrous sodium sulfate. Filtration and concentration gave crude 5-hydroxy-6-methoxynaphthalene (2.31 g, 46% yield) as a light brown solid which was used in the next step without further purification. LRMS (Electrospray): C₁₁H₁₀O₂, calc. 174; observed: 173 (M-H).


Step 2:



[0169] A mixture of 5-hydroxy-6-methoxynaphthalene (2.10 g, 12.1 mmole), cesium carbonate (19.7 g, 60.5 mmole), N,N-dimethylformamide (12 ml) and 2-bromopropane (3.50 ml, 36.9 mmole) was stirred in a 40°C oil bath overnight. The reaction mixture was filtered and the solid residue was rinsed thoroughly with ethyl ether. The filtrate and the washings were combined and concentrated in vacuo to remove most of the solvents. The brown residue was partitioned between water and ether and the layers were separated. The ether layer was washed with water. The combined aqueous layers were back


extracted with ether. The ether extracts were combined, washed with brine and dried over sodium sulfate. Filtration and concentration gave a crude which was purified by column chromatography (2.5 → 5% ethyl acetate/hexanes) to give 1-isopropoxy-6-methoxynaphthalene (2.23 g, 86% yield) as a light brown oil. ¹H NMR (CDCl₃) δ 8.17 (1H, d, phenyl), 7.05-7.38 (4H, m, phenyl), 6.72 (1H, dd, phenyl), 4.73 (1H, m, CH of iPr), 3.92 (3H, s, OCH₃), 1.42 (6H, d, 2 CH₃ of iPr).

Step 3:




[0170] To a refluxing solution of 1-isopropoxy-6-methoxynaphthalene (2.23 g, 10.3 mmole) in absolute ethanol (50 ml) was carefully added small pieces of sodium metal (3.6 g, 157 mmole) over 45 minutes. The mixture was refluxed for a further 120 minutes. The reaction was cooled to room temperature, quenched with water and acidified with concentrated hydrochloric acid. The mixture was concentrated in vacuo to remove most of the ethanol. The aqueous mixture was extracted three times with ether. The combined organic layers were washed with water and dried over sodium sulfate. Filtration and concentration gave a reddish oil which was dissolved in 1:1 ethanol/water (90 ml), then *p*-toluenesulfonic acid (200 mg) was added. The mixture was refluxed for 60 minutes. After cooling to room temperature, most of the ethanol was removed under reduced pressure. The aqueous mixture was extracted twice with ether and the combined organic layers were washed with water, saturated sodium chloride solution and dried over sodium sulfate. Filtration and concentration gave a reddish oil which was purified by column chromatography (8 → 15 % ethyl acetate/hexanes) to give 5-isopropoxy-β-tetralone (1.37 g, 65% yield) as a colorless oil. ¹H NMR (CDCl₃) δ 7.16 (1H, t, phenyl), 6.78 (1H, d, phenyl), 6.71 (1H, d, phenyl), 4.53 (1H, m, CH of iPr), 3.56 (2H, s, benzylic), 3.08 (2H, t, benzylic), 2.50 (2H, t), 1.37 (6H, d, 2 CH₃ of iPr).

Step 4:



[0171] A mixture of 5-isopropoxy- β -tetralone (1.37 g, 6.71 mmole), potassium cyanide (660 mg, 10.1 mmole), ammonium carbonate (3.87 g, 40.3 mmole), ethanol (44 ml) and water (9 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 42 hours. After cooling to room temperature, the slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave hydantoin Y (1.64 g, 89%).


Step 5:



[0172] A mixture of hydantoin Y (1.64 g, 5.98 mmole), Ba(OH)₂ · H₂O (5.66 g, 29.9 mmole) in water (25 ml) in a sealed, thick walled pressure flask was heated in a 100°C oil bath for 70 hours. The reaction mixture was cooled to room temperature, neutralized to ~ pH 7 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for one hour and cooled to room temperature. Basified with 1N sodium hydroxide solution and the white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 75 ml. Neutralization with concentrated hydrochloric acid solution gave white precipitate

which were filtered, washed with water and air-dried to give racemic 5-isopropoxy-2-aminotetraline-2-carboxylic acid (3.48 g, wet and containing inorganic salt, > quantitative yield). LRMS (Electrospray): $C_{14}H_{19}NO_3$, calc. 249; observed: 248 (M-H).

Step 6:




[0173] A mixture of racemic 5-isopropoxy-2-aminotetraline-2-carboxylic acid (3.48 g, 5.98 mmole theoretical), triethylamine (1.10 ml, 7.89 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 2.62 g, 7.77 mmole) in acetonitrile (30 ml) and water (30 ml) was stirred at room temperature for one day. TLC analysis of the reaction indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (500 mg) was added and the mixture was stirred at room temperature for another day. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted three times with methylene chloride. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 1 → 2 → 5 → 8% methanol/methylene chloride) to give racemic Fmoc-5-isopropoxy-2-aminotetraline-2-carboxylic acid (0.50 g, 18% yield over 2 steps) as a white solid. HRMS (FAB): $C_{29}H_{30}NO_5$ (M+H) calc. 472.2124; observed: 472.2117.

EXAMPLE 14


Preparation of Fmoc-(D,L)-5-methyl-2-aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-MeAtc-OH**)

Step 1:




[0174] A mixture of 2-methylhydrocinnamic acid (3.0 g, 18.3 mmole), oxalyl chloride (3.19 ml, 36.6 mmole) and methylene chloride (30 ml) was cooled in an ice bath and N,N-dimethylformamide (0.14 ml, 1.81 mmole) was added dropwise. The mixture was stirred at room temperature overnight. Concentration in vacuo gave 3-(2-methylphenyl)propanoyl chloride which was taken up in methylene chloride and used in the next step as a crude.

Step 2:




[0175] A solution of the above acid chloride (crude, 18.3 mmole) in methylene chloride was slowly added to a solution of diazomethane (generated from 11.9 g of 1-methyl-3-nitro-1-nitrosoguanidine) in ether (80 ml) cooled in an ice bath. The mixture was then warmed up to room temperature and stirred overnight. The mixture was concentrated in vacuo and purified by column chromatography (10 → 20% ethyl acetate/hexanes) to give 1-diazo-4-(2-methylphenyl)butan-2-one (2.08 g, 60% over 2 steps) as a bright yellow oil.

Step 3:




[0176] To a mixture of rhodium (II) acetate dimer (24 mg, 0.109 mmole) in methylene chloride (200 ml) under reflux was slowly added a solution of 1-diazo-4-(2-methylphenyl)butan-2-one (2.08 g, 11.1 mmole) in methylene chloride (50 ml) over 180 minutes. After the addition was complete, the mixture was refluxed for an extra twenty minutes. The mixture was cooled to room temperature, trifluoroacetic acid (2.40 ml) was added and the mixture was stirred at room temperature for an hour. The reaction was quenched with saturated sodium bicarbonate solution. The layers were separated and the methylene chloride layer was washed once more with saturated sodium bicarbonate solution. The combined aqueous layers were back-extracted with methylene chloride. The combined organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo to give a crude brown oil. Purification by column chromatography (15% ethyl acetate/hexanes) gave 5-methyl- β -tetralone (1.48 g, 84% yield) as a light brown oil. ^1H NMR (CDCl_3) δ 6.90-7.20 (3H, m, phenyl), 3.58 (2H, s, benzylic), 3.03 (2H, t, benzylic), 2.55 (2H, t), 2.34 (3H, s, CH_3).

Step 4:




[0177] A mixture of 5-methyl- β -tetralone (1.48 g, 9.24 mmole), potassium cyanide (902 mg, 13.9 mmole), ammonium carbonate (5.33 g, 55.5 mmole), ethanol (45 ml) and water (9 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 3 days. After cooling to room temperature, the slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave crude hydantoin **Z** (1.81 g, 85% yield) as a beige solid. ^1H NMR ($\text{DMSO}-d_6$) δ 10.66 (1H, broad s, NH), 8.22 (1H, broad s, NH), 6.85-7.05 (3H, m, phenyl), 2.17 (3H, s, CH_3).

Step 5:



[0178] A mixture of hydantoin **Z** (1.80 g, 7.82 mmole), Ba(OH)₂ · H₂O (7.40 g, 39.1 mmole) in water (28 ml) in a sealed, thick walled pressure flask was heated in a 125°C oil bath for 88 hours. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for an hour and cooled to room temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 50 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitate which were filtered, washed with water and air-dried to give racemic 5-methyl-2-aminotetraline-2-carboxylic acid (1.05 g, 65% yield) as a beige solid. LRMS (Electrospray): C₁₂H₁₅NO₂, calc. 205; observed: 206 (M+H).

Step 6:




[0179] A mixture of racemic 5-methyl-2-aminotetraline-2-carboxylic acid (1.05 g, 5.12 mmole), triethylamine (0.93 ml, 6.67 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 2.24 g, 6.64 mmole) in acetonitrile (30 ml) and water (30 ml) was stirred at room temperature for 2 days. TLC analysis of the reaction indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (520 mg) was added and the mixture was stirred at room temperature for another 24 hours. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified

to pH ~3 with 10% aqueous citric acid solution, and the white emulsion was extracted twice with methylene chloride. The combined organic layers were washed with water, brine and dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 2 → 5 → 8% methanol/methylene chloride) to give racemic Fmoc-5-methyl-2-aminotetraline-2-carboxylic acid (1.62 g, 74% yield) as an light brown solid. HRMS (FAB): C₂₇H₂₆NO₄ (M+H) calc. 428.1862; observed: 428.1844.

EXAMPLE 15


Preparation of Fmoc-(D,L)-5-ethyl-2 aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-EtAtc-OH**)

Step 1:




[0180] A mixture of 3-(2-ethylphenyl)propanoic acid (prepared in 3 steps from 1-ethyl-2-iodobenzene, 4.24 g, 23.8 mmole), thionyl chloride (9.50 ml, 130 mmole) and toluene (100 ml) was refluxed for 2 hours. Concentration in vacuo gave 3-(2-ethylphenyl)propanoyl chloride which was taken up in methylene chloride and used in the next step as a crude.

Step 2:




[0181] A solution of 3-(2-ethylphenyl)propanoyl chloride (crude, 23.8 mmole) in methylene chloride was slowly added to a solution of diazomethane (generated from 15.6 g of 1-methyl-3-nitro-1-nitrosoguanidine) in ether (100 ml) cooled in an ice bath. The mixture was then warmed up to room temperature and stirred overnight. The mixture was concentrated in vacuo and purified by column chromatography (10 → 20% ethyl acetate/hexanes) to give 1-diazo-4-(2-ethylphenyl)butan-2-one (3.47 g, 72% over 2 steps). ¹H NMR (CDCl₃) δ 7.1-7.25 (4H, m, phenyl), 5.21 (1H, broad s, diazo), 2.97 (2H, m, CH₂ of ethyl), 1.20 (3H, t, CH₃).

Step 3:




[0182] To a mixture of rhodium (II) acetate dimer (38 mg, 0.172 mmole) in methylene chloride (300 ml) under reflux was slowly added a solution of 1-diazo-4-(2-ethylphenyl)butan-2-one (3.47 g, 17.2 mmole) in methylene chloride (50 ml) over 90 minutes. After the addition was complete, the mixture was refluxed for an extra twenty minutes. The mixture was cooled to room temperature, trifluoroacetic acid (3.75 ml) was added and the mixture was stirred at room temperature for an hour. The reaction was quenched with saturated sodium bicarbonate solution. The layers were separated and the methylene chloride layer was washed once more with saturated sodium bicarbonate solution. The combined aqueous layers were back-extracted with methylene chloride. The combined organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo to give crude 5-ethyl-β-tetralone (3.09 g, > quantitative yield) as a reddish-brown oil. The compound was used in the next step without further purification. ¹H NMR (CDCl₃) δ 6.9-7.2 (3H, m, phenyl), 3.58 (2H, s, benzylic), 3.08 (2H, s, benzylic), 2.70 (2H, q, CH₂ of ethyl), 2.52 (2H, t, benzylic), 1.20 (3H, t, CH₃ of ethyl).

Step 4:



[0183] A mixture of 5-ethyl- β -tetralone (3.09 g, 17.7 mmole), potassium cyanide (1.73 g, 26.6 mmole), ammonium carbonate (10.2 g, 106 mmole), ethanol (80 ml) and water (16 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 48 hours. After cooling to room temperature, the white slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave hydantoin AA (3.85 g, 92% yield over 2 steps) as a light beige solid. ^1H NMR (DMSO- d_6) δ 10.67 (1H, broad s, NH), 8.26 (1H, broad s, NH), 6.8-7.1 (3H, m, phenyl), 1.13 (3H, t, CH_3). LRMS (Electrospray): $\text{C}_{14}\text{H}_{16}\text{N}_2\text{O}_2$, calc. 244; observed: 243 (M-H).


Step 5:



[0184] A mixture of hydantoin AA (1.00 g, 4.09 mmole), $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ (4.00 g, 21.1 mmole) in water (20 ml) in a sealed, thick walled pressure flask was heated in a 125°C oil bath for 48 hours. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for two hours and cooled to room temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 50 ml. Neutralization with concentrated

ammonium hydroxide solution gave white precipitate which were filtered, washed with water and dried in vacuo overnight to give racemic 5-ethyl-2-aminotetraline-2-carboxylic acid (796 mg, 89% yield). LRMS (Electrospray): $C_{13}H_{17}NO_2$, calc. 219; observed: 220 (M+H).

Step 6:




[0185] A mixture of racemic 5-ethyl-2-aminotetraline-2-carboxylic acid (765 mg, 3.49 mmole), triethylamine (1.0 ml, 7.17 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 1.79 g, 5.31 mmole) in acetonitrile (40 ml) and water (40 ml) was stirred at room temperature for 2 days. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted twice with methylene chloride, twice with ethyl acetate. The methylene chloride extracts were washed with water, brine and dried over magnesium sulfate. The ethyl acetate extracts were washed with water, brine and dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 2 → 5 → 8% methanol/methylene chloride) to give racemic Fmoc-5-ethyl-2-aminotetraline-2-carboxylic acid (330 mg, 21% yield) as a white solid. HRMS (FAB): $C_{28}H_{28}NO_4$ (M+H) calc. 442.2018; observed: 442.2010.

EXAMPLE 16


Preparation of Fmoc-(D,L)-5-isopropyl-2-aminotetraline-2-carboxylic acid (**Fmoc-(D,L) 5-iPrAtc-OH**)

Step 1:




[0186] A mixture of 3-(2-isopropylphenyl)propanoic acid (prepared in 3 steps from 1-isopropyl-2-iodobenzene, 2.01 g, 10.5 mmole), thionyl chloride (4.30 ml, 59.0 mmole) and toluene (40 ml) was refluxed for 2 hours. Concentration in vacuo gave 3-(2-isopropylphenyl)propanoyl chloride which was taken up in methylene chloride and used in the next step as a crude.

Step 2:




[0187] A solution of 3-(2-isopropylphenyl)propanoyl chloride (crude, 10.5 mmole) in methylene chloride was slowly added to a solution of diazomethane (generated from 6.95 g of 1-methyl-3-nitro-1-nitrosoguanidine) in ether (50 ml) cooled in an ice bath. The mixture was then warmed up to room temperature and stirred overnight. The mixture was concentrated in vacuo and purified by column chromatography (20% ethyl acetate/hexanes) to give 1-diazo-4-(2-isopropylphenyl)butan-2-one (1.87 g, 82% over 2 steps) as a bright yellow oil. ¹H NMR (CDCl₃) δ 7.10-7.30 (4H, m, phenyl), 5.21 (1H, broad s, diazo), 3.15 (1H, m, CH of iPr), 3.00 (2H, t, benzylic), 2.57 (2H, m), 1.24 (6H, d, 2 CH₃ of iPr).

Step 3:



[0188] To a mixture of rhodium (II) acetate dimer (20 mg, 0.091 mmole) in methylene chloride (160 ml) under reflux was slowly added a solution of 1-diazo-4-(2-bromophenyl)butan-2-one (1.87 g, 8.65 mmole) in methylene chloride (25 ml) over 60 minutes. After the addition was complete, the mixture was refluxed for an extra fifteen minutes. The mixture was cooled to room temperature, trifluoroacetic acid (1.90 ml) was added and the mixture was stirred at room temperature for 45 minutes. The reaction was quenched with saturated sodium bicarbonate solution. The layers were separated and the methylene chloride layer was washed once more with saturated sodium bicarbonate solution. The combined aqueous layers were back-extracted with methylene chloride. The combined organic layers were dried over magnesium sulfate, filtered and concentrated in vacuo to give a crude brown oil. Purification by column chromatography (5% ethyl acetate/hexanes) gave 5-isopropyl-β-tetralone (1.57 g, 96% yield) as a light yellow oil. ¹H NMR (CDCl₃) δ 6.93-7.22 (3H, m, phenyl), 3.59 (2H, s, benzylic), 3.24 (1H, m, CH of iPr), 3.12 (2H, t, benzylic), 2.52 (2H, t), 1.27 (6H, d, 2 CH₃ of iPr).


Step 4:



[0189] A mixture of 5-isopropyl-β-tetralone (1.57 g, 8.34 mmole), potassium cyanide (0.82 g, 12.6 mmole), ammonium carbonate (4.81 g, 50.1 mmole), ethanol (40 ml) and water (10 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 48 hours.

After cooling to room temperature, the brown slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration followed by air-drying gave crude hydantoin **BB** as a beige solid which was used in the next step without further purification. ^1H NMR (DMSO-d_6) δ 10.69 (1H, broad s, NH), 8.30 (1H, broad s, NH), 6.85-7.32 (3H, m, phenyl), 1.15 (6H, t, CH_3). LRMS (Electrospray): $\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_2$, calc. 258; observed: 539 (2M+Na).

Step 5:



[0190] A mixture of hydantoin **BB** (crude, 8.34 mmole theoretical), $\text{Ba}(\text{OH})_2 \cdot \text{H}_2\text{O}$ (7.90 g, 41.7 mmole) in water (40 ml) in a sealed, thick walled pressure flask was heated in a 125°C oil bath for 38 hours. The reaction mixture was cooled to room temperature, acidified to $\sim \text{pH } 3$ using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for two hours and cooled to room temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 50 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitate which were filtered, washed with water and dried in vacuo overnight to give racemic 5-isopropyl-2-aminotetraline-2-carboxylic acid (1.23 g, 63% yield over 2 steps) as a beige solid. LRMS (Electrospray): $\text{C}_{14}\text{H}_{19}\text{NO}_2$, calc. 233; observed: 232 (M-H).

Step 6:




[0191] A mixture of racemic 5-isopropyl-2-aminotetraline-2-carboxylic acid (250 mg, 1.07 mmole), triethylamine (1.2 ml, 8.61 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 2.70 g, 8.00 mmole) in acetonitrile (30 ml) and water (30 ml) was stirred at room temperature for 2 days. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion was extracted with ethyl acetate. The organic layer was washed with water, brine and dried over sodium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 2 → 5 → 8% methanol/methylene chloride) to give racemic Fmoc-5-isopropyl-2-aminotetraline-2-carboxylic acid (208 mg, 43% yield) as an off-white foam. HRMS (FAB): $C_{29}H_{30}NO_4$ (M+H) calc. 456.2175; observed: 456.2184.

EXAMPLE 17

Preparation of Fmoc-4-amino-1-phenylpiperidine-4-carboxylic acid (**Fmoc-Appc-OH**)


Step 1:



[0192] To a solution of iodobenzene (6.37 g, 31.2 mmole), 1,4-dioxaspiro[4.5]decane (10.32 g, 9.3 mL, 72.2 mmole, 2.3 equiv) and sodium tert-butoxide (8.0 g, 83.3 mmole, 2.7 equiv) in dry dioxane (120 mL) were added tris(dibenzylideneacetone)dipalladium(0) (91 mg, 0.1 mmol) and tri-*o*-tolylphosphine


(180 mg, 0.591 mmol). The reaction was heated at 90 °C for 26 hrs. The resulting reaction mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give a brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product **CC** as a slightly yellow solid (6.08 g, 89%). ¹H NMR (CDCl₃), 7.25 (ddt, 2H), 6.95 (dd, 2H), 6.84 (t, 1H), 4.00 (s, 4H), 3.32 (t, 4H) and 1.84 (t, 4H); MS (electrospray) *m/e* 220 (M+H), Calcd for C₁₃H₁₇NO₂, 219.

Step 2:




[0193] To a solution of the ketal **CC** (3.22 g, 15.16 mmol) in acetone (100 mL) was added 6N hydrochloric acid (50 mL) and the reaction was heated at reflux overnight. The resulting reaction mixture was concentrated to remove solvent. The residue was taken up in EtOAc and neutralized with aqueous 6N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 80/20→60/40) to give the product **DD** as a yellow oil (2.58 g, 97%). MS (electrospray) *m/e* 176 (M+H), Calcd for C₁₁H₁₃NO, 175.

Step 3:



[0194] To a solution of the ketone **DD** (2.53 g, 14.46 mmol) in ethanol (75 mL) and water (25 mL) in a glass pressure bottle, were added ammonium carbonate (12.9 g, 134.3 mmole, 9 equiv.) and potassium cyanide (2.11 g, 32.5 mmol, 2 equiv.). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was concentrated in vacuo and the residue was treated with water, extracted with EtOAc (4x). The combined organic extracts were washed with water, dried over anhydrous Na₂SO₄ and concentrated to give the spectroscopically pure hydantoin **EE** as a white solid (3.36 g, 95% yield). MS (electrospray) *m/e* 246 (M+H), Calcd for C₁₃H₁₅N₃O₂, 245.

Step 4:



[0195] The hydantoin **EE** (3.36 g) was suspended in aqueous NaOH (6N, 100 mL) and heated at 130 °C for 2-3 days. Upon completion (by HPLC) of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH ~6). The resulting slurry was filtered, washed with water and dried to give 4-amino-1-phenylpiperidine-4-carboxylic acid (APPC) as a white solid (5.26 g, >100 % yield, wet and contaminated with inorganic salt), which showed a single peak on HPLC and used directly for the next step. MS (electrospray) *m/e* 221 (M+H), Calcd for C₁₂H₁₆N₂O₂, 220.

Step 5:




[0196] The crude 4-amino-1-phenylpiperidine-4-carboxylic acid (APPC) from the last step was suspended in dioxane (80 mL) and aqueous 10% Na₂CO₃ (40 ml), treated with

Fmoc-Cl (5.3 g, 20.57 mmole, 1.5 equiv) and was stirred vigorously overnight. The reaction mixture was then concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na₂SO₄. Removal of the solvent gave the crude product which was purified on flash chromatography (hexane/EtOAc to CH₂Cl₂/MeOH) to give pure Fmoc-APPC (4.91 g, 81% overall yield for two steps). ¹H NMR(DMSO-d₆), 7.88 (d, 2H), 7.74 (d, 2H), 7.19-7.42 (m, 8H), 4.20-4.31 (m, 3H); HRMS m/z 465.1788, Calcd for C₂₇H₂₆N₂O₄Na, 465.1791.

EXAMPLE 18


Preparation of Fmoc-4-amino-1-(4-methylphenyl)piperidine-4-carboxylic acid (**Fmoc-4-MeAppc-OH**)

Step 1:




[0197] To a solution of 4-iodotoluene (2.12 g, 9.7 mmol), 1,4-dioxaspiro[4.5]decane (2.8 mL, 3.12 g, 21.82 mmol, 2.2 equiv) and sodium tert-butoxide (2.6 g, 27.08 mmol, 2.8 equiv) in dry dioxane (40 mL) were added tris(dibenzylideneacetone)dipalladium (0) (44.4 mg, 0.0485 mmol) and tri-*o*-tolylphosphine (59.0 mg, 0.194 mmol). The reaction was heated at 90 °C for 26 hrs. The resulting reaction mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product **FF** as a slightly yellow solid (1.937 g, 85%). ¹H NMR (CDCl₃), 7.06 (d, 2H), 6.87 (d, 2H), 3.99 (s, 4H), 3.26 (t, 4H), 2.26 (s, 3H) and 1.85 (t, 4H).

Step 2:




[0198] To a solution of the ketal **FF** (1.58 g, 6.79 mmol) in acetone (50 mL) was added 6N hydrochloric acid (25 mL) and the reaction was heated at reflux overnight. The resulting reaction mixture was concentrated to remove solvent. The residue was taken up in EtOAc and neutralized with aqueous 6N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 90/10→70/30) to give the product **GG** as a yellow oil (1.27 g, 98%). MS (electrospray) *m/e* 190 (M+H), Calcd for C₁₂H₁₅NO, 189.

Step 3:




[0199] To a solution of the ketone **GG** (1.17 g, 6.18 mmol) in ethanol (60 mL) and water (20 mL) in a glass pressure bottle, were added ammonium carbonate (4.74 g, 49.44 mmole, 8 equiv.) and potassium cyanide (1.01 g, 15.54 mmol, 2.5 equiv.). The mixture was heated at 90 °C for 22 hrs. The cooled reaction mixture was concentrated in vacuo and the residue was treated with water, extracted with EtOAc (4x). The combined organic extracts were washed with water, dried over anhydrous Na₂SO₄ and concentrated to give the spectroscopically pure hydantoin **HH** as a white solid (1.554 g, 97% yield). MS (electrospray) *m/e* 260 (M+H), Calcd for C₁₄H₁₇N₃O₂, 259.

Step 4:



[0200] The hydantoin **HH** (1.502 g) was suspended in aqueous NaOH (6N, 40 mL) and heated at 130 °C for 4 days. Upon completion (by HPLC) of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH ~6). The resulting slurry was filtered, washed with water and dried to give 4-amino-1-(4-methylphenyl)piperidine-4-carboxylic acid (4-MeAPPC) as a white solid (2.10 g, >100 % yield, wet and contaminated with inorganic salt), which showed a single peak on HPLC and used directly in the next step. MS (electrospray) *m/e* 235 (M+H), Calcd for C₁₃H₁₈N₂O₂, 234.

Step 5:




[0201] The crude 4-amino-1-(4-methylphenyl)piperidine-4-carboxylic acid (4-MeAPPC) from the last step was suspended in dioxane (80 mL) and aqueous 10% Na₂CO₃ (40 ml), treated with Fmoc-Cl (2.2 g, 8.59 mmole, 1.5 equiv) and was stirred vigorously overnight. The reaction mixture was then concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na₂SO₄. Removal of the solvent gave the crude product which was purified on flash chromatography (hexane/EtOAc to CH₂Cl₂/MeOH) to give pure Fmoc-4-MeAPPC (2.16 g, 82% overall yield for two steps). ¹H NMR (DMSO-d₆): 7.88 (d, 2H), 7.72 (d, 2H), 7.39 (t, 2H), 7.30 (td, 2H), 6.99 (d, 2H), 6.82 (d, 2H), 2.18 (s, 3H); MS (electrospray) *m/e* 457 (M+H), Calcd for C₂₈H₂₈N₂O₄, 456.

EXAMPLE 19


Preparation of Fmoc-4-amino-1-(4-chlorophenyl)piperidine-4-carboxylic acid (**Fmoc-4-ClAppc-OH**)

Step 1:



[0202] To a solution of 1-chloro-4-iodobenzene (2.38 g, 10.0 mmole), 1,4-dioxaspiro[3.5]nonan-8-amine (3.1 mL, 3.44 g, 24.0 mmole, 2.4 equiv) and sodium tert-butoxide (2.68 g, 28.0 mmole, 2.8 equiv) in dry dioxane (40 mL) were added tris(dibenzylideneacetone)dipalladium(0) (45.5 mg, 0.0497 mmol) and tri-*o*-tolylphosphine (61 mg, 0.20 mmol). The reaction was heated at 90 °C for 9 hrs. The resulting reaction mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give a brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product **II** as a slightly yellow solid (2.17 g, 86%). ¹H NMR(CDCl₃), 7.18 (dt, 2H), 6.85 (dt, 2H), 3.98 (s, 4H), 3.28 (t, 4H) and 1.82 (t, 4H).


Step 2:



[0203] To a solution of the ketal **II** (2.123 g, 8.39 mmole) in acetone (75 mL) was added 6N hydrochloric acid (30 mL) and the reaction was heated at reflux overnight. The resulting reaction mixture was concentrated to remove solvent. The residue was taken up in EtOAc and neutralized with aqueous 6N NaOH solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified


on flash chromatography (hexane/EtOAc, 95/5→70/30) to give the product **JJ** as a yellow solid (1.515 g, 86%). MS (electrospray) m/e 210 (M+H), Calcd for $C_{11}H_{12}ClNO$, 209.

Step 3:



[0204] To a solution of the ketone **JJ** (1.465 g, 6.986 mmole) in ethanol (75 mL) and water (25 mL) in a glass pressure bottle, were added ammonium carbonate (5.36 g, 55.88 mmole, 8 equiv.) and potassium cyanide (1.135 g, 17.46 mmol, 2.5 equiv.). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was concentrated in vacuo and the residue was treated with water, extracted with EtOAc (4x). The combined organic extracts were washed with water, dried over anhydrous Na_2SO_4 and concentrated to give the spectroscopically pure hydantoin **KK** as a white solid (1.817 g, 93% yield). MS (electrospray) m/e 280 (M+H), Calcd for $C_{13}H_{14}ClN_3O_2$, 279.

Step 4:



[0205] The hydantoin **KK** (1.768 g) was suspended in aqueous NaOH (6N, 50 mL) and heated at 130 °C for 4 days. Upon the completion (by HPLC) of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH ~6). The resulting slurry was filtered, washed with water and dried to give 4-amino-1-(4-

chlorophenyl)piperidine-4-carboxylic acid (4-ClAPPC) as a white solid (2.05 g, >100 % yield, wet and contaminated with inorganic salt), which showed a single peak on HPLC and used directly for the next step. MS (electrospray) m/e 253 (M-H), Calcd for $C_{12}H_{15}ClN_2O_2$, 254.

Step 5:




[0206] The crude 4-amino-1-(4-chlorophenyl)piperidine-4-carboxylic acid (4-ClAPPC) from the last step was suspended in dioxane (100 mL) and aqueous 10% Na_2CO_3 (50 ml), treated with Fmoc-Cl (2.0 g, 7.75 mmole, 1.2 equiv) and was stirred vigorously overnight. The reaction mixture was then concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na_2SO_4 . Removal of the solvent gave the crude product which was purified on flash chromatography (hexane/EtOAc to $CH_2Cl_2/MeOH$) to give pure Fmoc-4-ClAPPC (1.18 g, 81% overall yield for two steps). 1H NMR ($DMSO-d_6$): 7.87 (d, 2H), 7.71 (d, 2H), 7.39 (td, 2H), 7.30 (td, 2H), 7.20 (d, 2H), 6.92 (d, 2H), 3.44 (d, 2H), 2.93 (t, 2H); MS (electrospray) m/e 477 (M+H), Calcd for $C_{27}H_{25}N_2O_4$, 476.

EXAMPLE 20


Preparation of Fmoc-4-amino-1-(4-phenoxyphenyl)piperidine-4-carboxylic acid (Fmoc-4-PhOAppc-OH)

Step 1:




[0207] To a solution of 1-iodo-4-phenoxybenzene (3.15 g, 10.6 mmol), 1,4-dioxaspiro[3.5]nonan-8-amine (4.5 mL, 3.66 g, 25.6 mmole, 2.4 equiv) and sodium tert-butoxide (2.85 g, 29.7 mmol, 2.8 equiv) in dry dioxane (40 mL) were added tris (dibenzylideneacetone) dipalladium (0) (48.5 mg, 0.053 mmol) and tri-*o*-tolylphosphine (64 mg, 0.4 mmol). The reaction was heated at 90 °C for 9 hrs. The resulting reaction mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give a brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 80/20) to provide the pure product **LL** as a slightly yellow solid (2.805, 85%). ¹H NMR (CDCl₃), 7.26-7.32 (m, 2H), 7.03 (t, 1H), 6.92-6.97 (m, 6H), 4.00 (s, 4H), 3.26 (t, 4H), 1.86 (t, 4H).

Step 2:




[0208] To a solution of the ketal **LL** (2.755 g, 8.86 mmol) in acetone (90 mL) was added 6N hydrochloric acid (45 mL) and the reaction was heated at reflux overnight. The resulting reaction mixture was concentrated to remove solvent. The residue was diluted with EtOAc and neutralized with aqueous 6N NaOH. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated to give the crude product which was purified on flash chromatography (hexane/EtOAc, 90/10 to 70/30) to give the product **MM** as a yellow oil (2.21 g, 93%). MS (electrospray) *m/e* 268 (M+H), Calcd for C₁₇H₁₇ClNO₂, 267.

Step 3:




[0209] To a solution of the ketone **MM** (2.01 g, 7.52 mmol) in ethanol (80 mL) and water (25 mL) in a glass pressure bottle, were added ammonium carbonate (5.78 g, 60.0 mmol, 8 equiv.) and potassium cyanide (1.22 g, 18.80 mmol, 2.5 equiv.). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was concentrated in vacuo and the residue was treated with water, extracted with EtOAc (4x). The combined organic extracts were washed with water, dried over anhydrous Na₂SO₄ and concentrated to give the spectroscopically pure hydantoin **NN** as a white solid (2.34 g, 95% yield). MS (electrospray) *m/e* 338 (M+H), Calcd for C₁₉H₁₉N₃O₃, 337.

Step 4:



[0210] The hydantoin **NN** (2.28 g, 6.76 mmole) was suspended in aqueous NaOH (6N, 60 mL) and heated at 130 °C for 4 days. Upon completion (by HPLC) of the hydrolysis, the reaction mixture was neutralized with conc. HCl to slightly acidic (pH ~6). The resulting slurry was filtered, washed with water and dried to give 4-amino-1-(4-phenoxyphenyl)piperidine-4-carboxylic acid (4-PhOAPPC) as a white solid (2.53 g, >100 % yield, wet and contaminated with inorganic salt), which showed a single peak on HPLC and used directly for the next step. MS (electrospray) *m/e* 313 (M+H), Calcd for C₁₈H₂₀N₂O₃, 312.

Step 5:




[0211] The crude 4-amino-1-(4-phenoxyphenyl)piperidine-4-carboxylic acid (4-PhOAPPC) from the last step was suspended in dioxane (50 L) and aqueous 10 % Na₂CO₃ (50 ml) and stirred vigorously overnight. The reaction mixture was concentrated to remove dioxane, neutralized with 6N HCl to slightly acidic (pH 6) and extracted with EtOAc. The combined organic extracts were washed with brine and dried over Na₂SO₄. Removal of the solvent gave the crude product which was purified on flash chromatography (hexane/EtOAc to CH₂Cl₂/MeOH) to give pure Fmoc-4-PhOAPPC (2.18 g, 60% overall yield for two steps). ¹H NMR (DMSO-d₆): 7.87 (d, 2H), 7.72 (d, 2H), 7.38 (t, 2H), 7.30 (td, 4H), 7.02 (dt, 1H), 6.86-6.96 (m, 6H), 3.35 (m, 2H), 2.94 (t, 2H); MS (electrospray) *m/e* 535 (M+H), Calcd for C₃₃H₃₀N₂O₅, 534.

EXAMPLE 21

Preparation of Fmoc-4-amino-1-(2-methylphenyl)piperidine-4-carboxylic acid (Fmoc-2-MeAppc-OH)


Step 1:



[0212] To a solution of 2-iodotoluene (4.36 g, 2.5 mL, 20.0 mmol), 1,4-dioxaspiro[4.5]undecan-9-amine (6.88 g, 6.2 mL, 48.1 mmol, 2.4 equiv) and sodium tert-butoxide (5.3 g, 55.2 mmol, 2.8 equiv) in dry dioxane (80 mL) were added tris(dibenzylideneacetone)dipalladium(0) (91 mg, 0.1 mmol) and tri-*o*-tolylphosphine (122 mg, 0.4 mmol). The reaction was heated at 90 °C for 26 hrs. The resulting reaction


mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product **OO** as a slightly yellow solid (2.66 g, 57%). ¹H NMR (CDCl₃), 7.12-7.18 (m, 2H), 6.94-7.06 (m, 2H), 4.01 (s, 4H), 2.98 (t, 4H) and 1.88 (t, 4H).

Step 2:



[0213] To a solution of the ketal **OO** (2.66 g, 11.4 mmol) in acetone (70 mL) was added 6N hydrochloric acid (35 mL) and the reaction was heated at 85 °C overnight. The resulting reaction was concentrated to remove solvent. The residue was diluted with EtOAc and neutralized with aqueous NaOH (6N). The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 90/10 to 70/30) to give the product **PP** as a yellow oil (2.04 g, 95%). MS (electrospray) *m/e* 190 (M+H), Calcd for C₁₂H₁₅NO, 189.


Step 3:



[0214] To a solution of the ketone **PP** (1.54 g, 8.15 mmol) in ethanol (60 mL) and water (20 mL) in a glass pressure bottle, were added ammonium carbonate (4.69 g, 48.9 mmol, 6 equiv.) and potassium cyanide (800 g, 12.2 mmol, 1.5 equiv.). The mixture was heated at


80-90 °C for 18 hrs. The cooled reaction mixture was added to icy water (300 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **QQ** as a white solid (2.01 g, 95% yield). MS (electrospray) *m/e* 260 (M+H), Calcd for C₁₄H₁₇N₃O₂, 259.

Step 4:



[0215] To a suspension of the hydantoin **QQ** (1.07 g, 4.13 mmol) in dry THF (25 mL) were added di-*tert*-butyl dicarbonate (2.25 g, 10.32 mmol, 2.5 equiv), triethylamine (0.63 mL, 460 mg, 4.54 mmol, 1.1 equiv) and DMAP (36 mg, 0.29 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (300 mL), washed with 1N HCl (3x30 mL), saturated aqueous Na₂CO₃ (2x30 mL) and brine (2x30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→80/20) to give the pure bis-Boc hydantoin **RR** as a white solid (1.71 g, 90%). MS (electrospray) *m/e* 460 (M+H), Calcd for C₂₄H₃₃N₃O₆, 459.

Step 5:




[0216] The bis-Boc hydantoin **RR** (1.71g, 3.72 mmol) was dissolved in DME (23 mL) to give a clear solution. To this solution was added 1N NaOH (33 mL, 33 mmol) and the

reaction was stirred overnight at room temperature, giving a fairly clear mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 4-amino-1-(2-methylphenyl)piperidine-4-carboxylic acid (2-MeAPPC) was treated with 6N HCl to adjust the pH to 11-12. This solution (30 mL) was then diluted with 1,4-dioxane (30 mL) and treated with Fmoc-Cl (1.28 g, 4.96 mmol, 1.3 equiv) and stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove dioxane, neutralized with 1N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (hexane/EtOAc→CH₂Cl₂/MeOH) to give the pure product Fmoc-2-MeAPPC as a white solid (1.09 g, 64 % yield from the bis-Boc hydantoin **RR**). ¹H NMR (DMSO-d₆): 7.87 (d, 2H), 7.74 (d, 2H), 7.40 (td, 2H), 7.31 (td, 2H), 7.12 (m, 2H), 6.97 (d, 1H), 6.92 (td, 1H), 2.72-2.88 (m, 4H) and 2.22 (s, 3H); MS (electrospray) *m/e* 457 (M+H), Calcd for C₂₈H₂₈N₂O₄, 456.

EXAMPLE 22

Preparation of Fmoc-4-amino-1-(2-isopropylphenyl)piperidine-4-carboxylic acid (**Fmoc-2-iPrAppc-OH**)


Step 1:



[0217] To a solution of 1-iodo-2-*iso*-propylbenzene (10.0 g, 40.7 mmol), 1,4-dioxaspiro[4.5]undecane (12.0 mL, 13.3 g, 93.0 mmol, 2.3 equiv) and sodium *tert*-butoxide (10.0 g, 104.2 mmol, 2.6 equiv) in dry dioxane (160 mL) were added tris(dibenzylideneacetone)dipalladium(0) (180 mg, 0.197 mmol) and tri-*o*-tolylphosphine (244 mg, 0.80 mmol) and the reaction was heated at 90 °C for 26 hrs. The resulting reaction mixture was concentrated to remove solvent, treated with water and


extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give a brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5→75/25) to provide the pure product SS as a slightly yellow solid (3.61 g, 35% yield). MS *m/z* 262 (M+H), Calcd for C₁₆H₂₃NO₂, 261.

Step 2:



[0218] To a solution of the ketal SS (3.24 g, 12.4 mmol) in acetone (90 mL) was added 6N hydrochloric acid (45 mL) and the reaction was heated at reflux overnight. The resulting reaction mixture was concentrated to remove solvent and the residue was diluted with EtOAc, neutralized with aqueous NaOH (6N). The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 90/10→70/30) to give the product TT as a yellow oil (2.42 g, 89%). ¹H NMR (CDCl₃): 7.27 (m, 1H), 7.04-7.19 (m, 3H), 3.58 (m, 1H), 3.20 (t, 4H), 2.60 (t, 4H) and 1.25 (d, 6H); MS *m/z* 218 (M+H), Calcd for C₁₄H₁₉NO, 217.


Step 3:



[0219] To a solution of the ketone TT (2.30 g, 10.6 mmol) in ethanol (90 mL) and water (20 mL) in a glass pressure bottle, were added ammonium carbonate (8.1 g, 84.3 mmol, 8


equiv) and potassium cyanide (1.72 g, 26.5 mmol, 2.5 equiv). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was added to icy water (400 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin UU as a white solid (2.78 g, 91% yield). MS *m/z* 288 (M+H), Calcd for C₁₆H₂₁N₃O₂, 287.

Step 4:



[0220] To a suspension of the hydantoin UU (2.74 g, 9.54 mmol) in dry THF (100 mL) were added di-*tert*-butyl dicarbonate (5.2 g, 24.24 mmol, 2.5 equiv), triethylamine (1.5 mL, 1.07 g, 10.5 mmol, 1.1 equiv) and DMAP (46 mg, 0.29 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (300 mL), washed with brine (3x30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→80/20) to give the pure bis-Boc hydantoin VV as a white solid (4.39 g, 94% yield). MS *m/z* 488 (M+H), Calcd for C₂₆H₃₇N₃O₆, 487.

Step 5:




[0221] The bis-Boc hydantoin **VV** (2.34g, 4.8 mmol) was dissolved in DME (30 mL) to give a clear solution. To this solution was added 1N NaOH (45 mL, 45 mmol) and the reaction was stirred overnight at room temperature, giving a fairly clear mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 4-amino-1-(2-isopropylphenyl)piperidine-4-carboxylic acid (2-iPrAPPC) was treated with 6N HCl to adjust the pH to 11-12. This solution (~45 mL) was then diluted with 1,4-dioxane (45 mL) and treated with Fmoc-Cl (1.78 g, 6.89 mmol, 1.5 equiv) and stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove dioxane, neutralized with 1N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (hexane/EtOAc→CH₂Cl₂/MeOH) to give the pure product Fmoc-2-iPrAPPC as a white solid (1.46 g, 63 % yield from the bis-Boc hydantoin). HRMS m/z 507.2263, Calcd for C₃₀H₃₂N₂O₄Na, 507.2260.

EXAMPLE 23

Preparation of Fmoc-4-amino-1-(3-methylphenyl)piperidine-4-carboxylic acid (**Fmoc-3-MeAppc-OH**)


Step 1:



[0222] To a solution of 3-iodotoluene (4.36 g, 2.6 mL, 20.0 mmol), 1,4-dioxo-8-azaspiro [4.5] decane (6.88 g, 6.2 mL, 48.1 mmol, 2.4 equiv) and sodium tert-butoxide (5.3 g, 55.2 mmol, 2.8 equiv) in dry dioxane (80 mL) were added tris (dibenzylideneacetone) dipalladium (0) (91 mg, 0.1 mmol) and tri-*o*-tolylphosphine (122 mg, 0.4 mmol). The


reaction was heated at 90 °C for 26 hrs. The resulting reaction mixture was concentrated to remove solvent. The residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give a brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product **WW** as a slightly yellow solid (3.21 g, 69%).

Step 2:




[0223] To a solution of the ketal **WW** (1.25 g, 5.36 mmol) in acetone (20 mL) was added 6N hydrochloric acid (10 mL) and the reaction was heated at reflux overnight. The resulting reaction was concentrated to remove solvent. The residue was diluted with EtOAc and neutralized with aqueous NaOH (6N). The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 90/10 to 70/30) to give the product **XX** as a yellow oil (843 mg, 83% yield). MS *m/z* 190 (M+H), Calcd for C₁₂H₁₅NO, 189.

Step 3:



[0224] To a solution of the ketone **XX** (763 g, 4.03 mmol) in ethanol (45 mL) and water (15 mL) in a glass pressure bottle, were added ammonium carbonate (3.09 g, 32.21 mmol, 8 equiv) and potassium cyanide (675 mg, 10.38 mmol, 2.5 equiv). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was added to icy water (200 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **YY** as a white solid (930 mg, 89% yield). MS *m/z* 260 (M+H), Calcd for C₁₄H₁₇N₃O₂, 259.

Step 4:



[0225] To a suspension of the hydantoin **YY** (780 mg, 3.012 mmol) in dry THF (22 mL) were added di-*tert*-butyl dicarbonate (1.64 g, 7.52 mmol, 2.5 equiv), triethylamine (0.42 mL, 305 mg, 3.01 mmol, 1.0 equiv) and DMAP (20 mg, 0.164 mmol) in succession. About 5 minutes after the addition, the reaction turned into a clear yellow solution and was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (300 mL), washed with brine (3x30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→80/20) to give the pure bis-Boc hydantoin **ZZ** as a white solid (1.37 g, quantitative). HRMS *m/z* 482.2261 (M+Na), Calcd. for C₂₄H₃₃N₃O₆Na, 482.2267.

Step 5:




[0226] The bis-Boc hydantoin **ZZ** (1.29 g, 2.818 mmol) was dissolved in DME (20 mL) to give a clear solution. To this solution was added 1N NaOH (25 mL, 25 mmol) and the reaction was stirred overnight at room temperature, giving a fairly clear mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 4-amino-1-(3-methylphenyl)piperidine-4-carboxylic acid (3-MeAPPC) was treated with 6N HCl to adjust the pH to 11-12. This solution (30 mL) was then diluted with 1,4-dioxane (30 mL) and treated with Fmoc-Cl (1.46 mg, 5.65 mmol, 2.0 equiv) and stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove dioxane, neutralized with 1N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (hexane/EtOAc → CH₂Cl₂/MeOH) to give the pure product Fmoc-3-MeAPPC as a white solid (1.002 g, 78 % yield from the bis-Boc hydantoin). HRMS m/z 479.1940 (M+Na), Calcd. for C₂₈H₂₈N₂O₄Na, 479.1947.

EXAMPLE 24


Preparation of Fmoc-4-amino-1-(3-methoxyphenyl)piperidine-4-carboxylic acid (**Fmoc-3-MeOAppc-OH**)

Step 1:



[0227] To a solution of 3-iodoanisole (4.68 g, 2.4 mL, 20.0 mmol), 1,4-dioxaspiro [4.5] decane (6.2 mL, 6.88 g, 48.1 mmol, 2.4 equiv) and sodium tert-butoxide (5.3 g, 55.2 mmol, 2.8 equiv) in dry dioxane (80 mL) were added tris(dibenzylideneacetone)dipalladium(0) (91 mg, 0.1 mmol) and tri-*o*-tolylphosphine (122 mg, 0.4 mmol) and the reaction was heated at 90 °C for 26 hrs. The resulting reaction mixture was concentrated to remove solvent and the residue was treated with water and extracted with EtOAc. The combined organic extracts were combined, washed with brine, dried over Na₂SO₄ and concentrated to give brown oil. This crude product was purified on flash chromatography (hexane/EtOAc, 95/5 to 75/25) to provide the pure product AAA as a slightly yellow solid (3.10 g, 62% yield). MS *m/z* (M+H), 250 (M+H), Calcd for C₁₄H₁₉NO₃, 249.


Step 2:



[0228] To a solution of the ketal AAA (3.10 g, 12.45 mmol) in acetone (90 mL) was added 6N hydrochloric acid (45 mL) and the reaction was heated at reflux overnight. The resulting reaction was concentrated to remove solvent. The residue was diluted with EtOAc and neutralized with aqueous NaOH (6N). The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed


with brine, dried over Na₂SO₄ and concentrated. The crude product was purified on flash chromatography (hexane/EtOAc, 90/10 to 70/30) to give the product **BBB** as a yellow oil (2.53 g, 99% yield). ¹H NMR (CDCl₃): 7.20 (m, 1H), 6.58 (d, 1H), 6.39-6.56 (m, 2H), 3.80 (s, 3H), 3.59 (m, 4H) and 2.58 (m, 4H).

Step 3:



[0229] To a solution of the ketone **BBB** (1.81 g, 8.82 mmol) in ethanol (60 mL) and water (20 mL) in a glass pressure bottle, were added ammonium carbonate (6.77 g, 70.52 mmol, 8 equiv) and potassium cyanide (1.14g, 17.6 mmol, 2.0 equiv). The mixture was heated at 80-90 °C for 18 hrs. The cooled reaction mixture was added to icy water (200 ml) and stirred vigorously for 30 min. The resulting precipitate was suction filtered, washed thoroughly with water and dried to yield the hydantoin **CCC** as a white solid (2.23 g, 92% yield). MS *m/z* 276 (M+H), Calcd for C₁₄H₁₇N₃O₃, 275.


Step 4:



[0230] To a suspension of the hydantoin **CCC** (1.10 g, 4.00 mmol) in dry THF (50 mL) were added di-*tert*-butyl dicarbonate (2.18 g, 10.0 mmol, 2.5 equiv), triethylamine (0.62 mL, 445 mg, 4.4 mmol, 1.1 equiv) and DMAP (20 mg, 0.164 mmol) in succession. About 15 minutes after the addition, the reaction turned into a clear yellow solution and was stirred

overnight at room temperature. The reaction mixture was concentrated under reduced pressure to yield a solid that was then taken up in EtOAc (300 mL), washed with brine (3x30 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The crude light yellow product was purified through flash chromatography (hexane/EtOAc, 90/10→80/20) to give the pure bis-Boc hydantoin **DDD** as a white solid (1.90 g, quantitative). ¹H NMR (CDCl₃): 7.16 (t, 1H), 6.57 (d, 1H), 6.24 (s, 1H), 6.19 (d, 1H), 3.77 (s, 3H), 1.58 (s, 9H), 1.42 (s, 9H); MS *m/z* 476 (M+H), Calcd for C₂₄H₃₃N₃O₇, 475.

Step 5:




[0231] The bis-Boc hydantoin **DDD** (1.06 g, 2.23 mmol) was dissolved in DME (20 mL) to give a clear solution. To this solution was added 1N NaOH (20 mL, 20 mmol) and the reaction was stirred overnight at room temperature, giving a fairly clear mixture. HPLC showed completion of the reaction. The reaction mixture was concentrated under reduced pressure to remove DME and extracted with Et₂O. Without purification, the resulting aqueous layer containing 4-amino-1-(3-methoxyphenyl)piperidine-4-carboxylic acid (3-MeOAPPC) was treated with 6N HCl to adjust the pH to 11-12. This solution (35 mL) was then diluted with 1,4-dioxane (35 mL) and treated with Fmoc-Cl (755 mg, 2.93 mmol, 1.3 equiv) and stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure to remove dioxane, neutralized with 1N HCl and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated. The crude product was purified through flash chromatography (hexane/EtOAc→ CH₂Cl₂/MeOH) to give the pure product Fmoc-3-MeOAPPC as a white solid (668 mg, 63 % yield from the bis-Boc hydantoin **DDD**). ¹H NMR (CDCl₃): 7.83 (d, 2H), 7.72 (d, 2H), 7.41 (td, 2H), 7.34 (dt, 2H), 7.16 (t, 1H), 6.52 (d, 1H), 6.42 (s, 1H), 6.36 (d, 1H), 4.25 (m, 3H), 3.68 (s, 3H), 3.23-3.40 (m, 2H), 2.96 (t,

2H) and 1.86-2.18 (m, 4H). HRMS m/z 495.1901 ($M+Na$), Calcd. for $C_{28}H_{28}N_2O_5Na$, 495.1896.

EXAMPLE 25


Preparation of Fmoc-1-amino-4-cyclohexylcyclohexane-1-carboxylic acid (**Fmoc-Achc-OH**)

Step 1:




[0232] A mixture of 4-cyclohexylcyclohexanone (3.00 g, 16.6 mmole), potassium cyanide (1.63 g, 25.0 mmole), ammonium carbonate (9.59 g, 99.8 mmole), ethanol (75 ml) and water (15 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 15 hours. After cooling to room temperature, the white slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration and air-drying gave hydantoin **EEE** (6.10 g, still wet, >100% yield) as a white solid. 1H NMR (DMSO- d_6) δ 10.52 (1H, broad, NH), 8.43 (1H, broad s, NH), 0.80-1.80 (20H, m). LRMS (APCI): $C_{14}H_{22}N_2O_2$, calc. 250; observed: 249 ($M-H$), 251 ($M+H$).

Step 2:



[0233] A mixture of hydantoin **EEE** (1.39 g, 5.55 mmole) and 6N sodium hydroxide solution (50 ml) in a sealed, thick walled pressure flask was heated in a 130°C oil bath for 2 days. The reaction mixture was cooled in an ice bath, neutralized to ~ pH 7 using concentrated hydrochloric acid. The white slurry was filtered and the precipitates rinsed with water to give crude 1-amino-4-cyclohexylcyclohexane-1-carboxylic acid (48.3 g, wet and containing inorganic salts, >100% yield). LRMS (Electrospray): $C_{13}H_{23}NO_2$, calc. 225; observed: 226 (M+H).

Step 3:




[0234] A mixture of crude 1-amino-4-cyclohexylcyclohexane-1-carboxylic acid (48.3 g, 5.55 mmole theoretical), triethylamine (1.0 ml, 7.17 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 2.43 g, 7.20 mmole) in acetonitrile (75 ml) and water (75 ml) was stirred at room temperature for 24 hours. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted three times with methylene chloride. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 1 → 5 → 8% methanol/methylene chloride) to give Fmoc-1-amino-4-*trans*-cyclohexylcyclohexane-1-carboxylic acid (250 mg, 10% yield for two steps). HRMS (FAB): $C_{28}H_{34}NO_4$ (M+H) calc. 448.2488; observed: 448.2497.

EXAMPLE 26


Preparation of Fmoc-1-amino-4,4-diphenylcyclohexane-1-carboxylic acid (**Fmoc-Adpc-OH**)

Step 1:



[0235] A mixture of 4,4-diphenylcyclohexanone (prepared by hydrogenation of 4,4-diphenylcyclohexenone according to the procedures of Freeman, P.K. *et.al. J. Org. Chem.* **1989**, *54*, 782-789) (1.55 g, 6.19 mmole), potassium cyanide (0.65 g, 9.97 mmole), ammonium carbonate (3.60 g, 37.5 mmole), ethanol (48 ml) and water (12 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 24 hours. After cooling to room temperature, the white slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration and air-drying gave hydantoin **FFF** (1.89 g, 95% yield) as a white solid. ¹H NMR (DMSO-d₆) δ 10.57 (1H, broad, NH), 8.59 (1H, broad s, NH), 7.00-7.50 (10H, m, phenyl). LRMS (Electrospray): C₂₀H₂₀N₂O₂, calc. 320; observed: 319 (M-H).


Step 2:



[0236] A mixture of hydantoin **FFF** (1.88 g, 5.87 mmole), barium hydroxide monohydrate (5.60 g, 29.6 mmole) and water (100 ml, too dilute!) in a sealed, thick walled pressure flask was heated in a 105°C oil bath for 2 days. More barium hydroxide monohydrate (5.60 g, 29.6 mmole) was added and the mixture was heated in a 105°C oil bath for another 24 hours. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for two hours and cooled to room temperature. The white

suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 30 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitates which were filtered, washed with water and dried in vacuo overnight to give crude 1-amino-4,4-diphenylcyclohexane-1-carboxylic acid (0.52 g, 30% yield) as a white solid. LRMS (Electrospray): $C_{19}H_{21}NO_2$, calc. 295; observed: 294 (M-H), 296 (M+H).

Step 3:




[0237] A mixture of crude 1-amino-4,4-diphenylcyclohexane-1-carboxylic acid (510 mg, 1.73 mmole), triethylamine (0.37 ml, 2.65 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 880 mg, 2.61 mmole) in acetonitrile (25 ml) and water (25 ml) was stirred at room temperature overnight. TLC analysis of the reaction indicated the presence of starting material amino acid. 9-fluorenylmethyl succinimidyl carbonate (200 mg) and acetonitrile (5 ml) were added and the mixture was stirred at room temperature for another 24 hours. The reaction mixture was concentrated in vacuo to remove most of the acetonitrile, acidified to pH ~3 with 10% aqueous citric acid solution, and the white emulsion extracted three times with ethyl acetate. The combined organic layers were washed with water, brine, dried over sodium sulfate. Filtration and concentration gave a crude oil which was purified by column chromatography (eluted with 1 → 4 → 8% methanol/methylene chloride) to give Fmoc-1-amino-4,4-diphenylcyclohexane-1-carboxylic acid (350 mg, 39% yield) as a white solid. HRMS (FAB): $C_{34}H_{32}NO_4$ (M+H) calc. 518.2331; observed: 518.231

EXAMPLE 27


Preparation of Fmoc-1-amino-4-*trans*-*t*-butylcyclohexane-1-carboxylic acid (**Fmoc-Abc-OH**)

Step 1:



[0238] A mixture of 4-*t*-butylcyclohexanone (2.00 g, 13.0 mmole), potassium cyanide (1.27 g, 19.5 mmole), ammonium carbonate (7.48 g, 77.8 mmole), ethanol (60 ml) and water (12 ml) in a sealed, thick walled pressure flask was heated in a 80°C oil bath for 15 hours. After cooling to room temperature, the white slurry was poured into ice-water and stirred at room temperature for a couple of hours. Filtration gave hydantoin **GGG** (2.78 g, 96% yield) as a white solid which was used in the next step as a crude. ¹H NMR (DMSO-*d*₆) δ 10.52 (1H, broad, NH), 8.50 (1H, broad s, NH), 0.81 (9H, s, *t*-Bu).


Step 2:



[0239] A mixture of hydantoin **GGG** (2.78 g, 12.4 mmole), barium hydroxide monohydrate (11.74 g, 62.0 mmole) and water (50 ml) in a sealed, thick walled pressure flask was heated in a 120°C oil bath for 2 days. The reaction mixture was cooled to room temperature, acidified to ~ pH 3 using 4N sulfuric acid while being stirred vigorously. The suspension was stirred in a boiling water bath for one hour and cooled to room

temperature. The white suspension was filtered and the precipitates rinsed with water. The combined filtrate and washings were concentrated in vacuo to ~ 30 ml. Neutralization with concentrated ammonium hydroxide solution gave white precipitates which were filtered, washed with water and dried in vacuo overnight to give 1-amino-4-*trans*-*t*-butylcyclohexane-1-carboxylic acid (2.10 g, 85% yield) as a white solid.

Step 3:




[0240] A mixture of crude 1-amino-4-*trans*-*t*-butylcyclohexyl-1-carboxylic acid (2.10 g, 10.54 mmole), 9-fluorenylmethyl succinimidyl carbonate (Fmoc-OSu, 6.33 g, 7.20 mmole) in dioxane (150 ml) and 10% sodium carbonate solution (120 ml) was stirred at room temperature for 24 hours. The reaction mixture was concentrated in vacuo to remove most of the dioxane, acidified to pH ~3 with 3N HCl, and the white emulsion extracted twice with methylene chloride. The combined organic layers were washed with water, brine, dried over magnesium sulfate. Filtration and concentration gave a crude which was purified by column chromatography (eluted with 1 → 4 → 5% methanol/methylene chloride) to give Fmoc-1-amino-4-*trans*-*t*-butylcyclohexane-1-carboxylic acid (1.42 g, 32% yield). HRMS (FAB): C₂₆H₃₂NO₄ (M+H) calc. 422.2331; observed: 422.23.

EXAMPLE 28

Preparation of 3*S*, 2*S*-Fmoc-(*L*)-beta methy (Nin-Mes) tryptophan, **Fmoc-(*L*)-β-Me(Nin-Mes)Trp-OH**)

Step 1:



[0241] To a solution of trans-3-indoleacrylic acid (15.0 g, 0.08 mole) in 350 mL of dry THF at -78 °C was slowly added 125 mL of 1.6 M n-BuLi in hexane. The resulting suspension was stirred at -78 °C for 1 hr. Then, a solution of 2-mesitylenesulfonyl chloride (21.9 g, 0.1 mole) in 50 mL of dry THF was slowly added. The mixture was warmed to rt, and stirred overnight. The mixture was poured to a saturated aqueous NH₄Cl solution. The layers were separated and the aqueous layer was extracted with EtOAc. The combined organic layer was dried over sodium sulfate. Removal of solvents gave 14.1 g of crude product **HHH** which was used for the next step without further purification. ¹H NMR analysis indicates that it contains 2.8 g of 2-mesitylenesulfonic acid. ¹H NMR (CD₃OD) δ 7.57 (s, 1H), 7.42 (d, 1H), 7.15-7.30 (m, 3H), 7.02 (s, 2H), 6.54 (d, 1H), 6.36 (d, 1H), 2.52 (s, 9H), 2.30 (s, 3H).


Step 2:



[0242] To a solution of N-2-mesitylenesulfonyl-trans-3-indoleacrylic acid (3.26 g, 8.8 mmole) in 140 mL of dry THF at -78 °C were added 3.7 mL (3 equiv.) of triethylamine and 2.17 mL (2 equiv.) of Me₃CCOCl. The resulting mixture was stirred at -78 °C for 15 min, and at 0 °C for 1.5 hr. The mixture was cooled down to -78 °C, to which was added 5.5 mL of 1.6M n-BuLi in hexane, then added a mixture of (R)-4-phenyl-2-oxazolidinone and n-BuLi in THF (prepared by adding 11 mL of 1.6M n-BuLi in hexane to a solution of (R)-4-phenyl-2-oxazolidinone (2.87 g, 17.6 mmole) in 70 mL of dry THF at -78 °C) through a cannula. The resulting mixture was stirred at -78 °C for 2 hr, and at rt overnight. The


reaction was quenched by NH_4Cl aqueous solution (100 mL). After removal of the organic solvents in vacuo, the aqueous residue was extracted with EtOAc. The combined organic layer was dried over sodium sulfate. Filtration and concentration gave a crude product which was purified by a flash chromatography (EtOAc/hexane, 1:4) to give the product **III** as light brown gum in 63% yield (2.86 g). LR-Electrospray: $\text{C}_{29}\text{H}_{26}\text{N}_2\text{O}_5\text{S}$, cal: 514 observed: m/z 515 ($\text{M}+\text{H}$).

Step 3:




[0243] To a mixture of $\text{CuBr} \cdot \text{Me}_2\text{S}$ (0.84 g, 4.08 mmole) and 5 mL of dimethyl sulfide in 10 mL of dry THF at -4°C was added 1.36 mL of 3M CH_3MgBr in ether. After stirring for 10 min, the above product (1.4 g, 2.72 mmole) in 8 mL of dry THF was added. The resulting mixture was stirred at -4°C for 1 hr, and at rt for 6 hr. After cooling to -78°C , to the mixture was added 1.45 g (8.16 mmole) of N-bromosuccinimide in 15 mL of dry THF. The mixture was stirred at -78°C for 30 min, and at rt overnight. The mixture was poured to 100 mL of brine, and extracted with EtOAc (2x100 mL). The organic layer was dried over sodium sulfate. Filtration and concentration gave a crude product which was purified by a flash chromatography (EtOAc/hexane, 1:4) to give the product **JJJ** as light brown gum in 46% yield (0.77 g). ^1H NMR (CDCl_3) δ 7.63 (d, 1H), 7.47 (s, 1H), 7.20-7.37 (m, 8H), 6.98 (s, 2H), 6.16 (d, 1H), 5.13 (dd, 1H), 4.49 (t, 1H), 4.17 (dd, 1H), 3.75 (dt, 1H), 2.54 (s, 9H), 2.31 (s, 3H), 1.59 (d, 3H).

Step 4:



[0244] The above bromide **JJJ** (0.72 g, 1.18 mmole) was mixed with tetra-n-butylammonium azide (1.68 g, 5.9 mmole) and sodium azide (77 mg, 1.18 mmole) in 10 mL of acetonitrile, and stirred at rt for 6 hr. The mixture poured to 100 mL of NH_4Cl aqueous solution and extracted with EtOAc (2x100 mL). The organic layer was dried over sodium sulfate. Filtration and concentration gave a crude product which was purified by a flash chromatography (EtOAc/ CH_2Cl_2 /hexane, 1:2:5) to give the product **KKK** as light brown gum in 82% yield (0.55 g). 1H NMR ($CDCl_3$) δ 7.65 (d, 1H), 7.59 (s, 1H), 7.18-7.29 (m, 8H), 6.90 (s, 2H), 5.54 (d, 1H), 5.50 (dd, 1H), 4.78 (t, 1H), 4.35 (dd, 1H), 3.62 (quintet, 1H), 2.43 (s, 9H), 2.28 (s, 3H), 1.28 (d, 3H).


Step 5:



[0245] To a mixture of the above azide **KKK** (0.55 g, 0.96 mmole), water (4 mL) and THF (12 mL) at 0 °C was added 0.65 mL of 30% H_2O_2 , then added 48 mg (2 equiv.) of LiOH in 1 mL water. The resulting mixture was stirred at 0 °C for 2 hr. The reaction was quenched with Na_2SO_3 (1 g) in 6 mL water. The mixture was stirred at rt for additional 30 min. After removal of the organic solvent, the aqueous solution was diluted with 10 mL of saturated $NaHCO_3$ solution and extracted with EtOAc (2x30 mL). Filtration and

concentration gave a crude product which was purified by a flash chromatography (HOAc/MeOH/EtOAc, 1:10:100) to give the product **LLL** as off-white solid in 83% yield (0.34 g). LR-Electrospray: $C_{21}H_{22}N_4O_4S$, cal: 426 observed: m/z 425 (M-H).

Step 6:



[0246] The above azido acid **LLL** (0.34 g, 0.8 mmole) was dissolved in 20 mL of methanol.

To the solution was added 170 mg of 10% Pd on carbon. The resulting mixture was stirred at rt under H_2 (balloon) for 3 hr. After filtration and concentration, the crude was dissolved in a mixed solvent of THF (12 mL) and water (4 mL). To the mixture were added $NaHCO_3$ (254 mg, 3 mmole) and Fmoc-OSu (540 mg, 1.6 mmole). The resulting mixture was stirred at rt for 18 hr. Diluted with 30 mL of saturated NH_4Cl solution and extracted with EtOAc (2x30 mL). Filtration and concentration gave a crude product which was purified by a flash chromatography (HOAc/MeOH/EtOAc, 1:10:100) to give the product 3*S*, 2*S*-Fmoc-(L)-beta methy (Nin-Mes) tryptophan as off-white solid in 50% yield (0.25 g). LR-Electrospray: $C_{36}H_{34}N_2O_6S$, cal: 622 observed: m/z 621 (M-H).

EXAMPLE 29


Preparation of Fmoc-Linker-BHA Resin

[0247] Benzhydrylamine copolystyrene-1% divinylbenzene cross-linked resin (10.0 g, 9.3 mequiv, 100-200 ASTM mesh, Advanced ChemTech) was swelled in 100 mL CH_2Cl_2 , filtered and washed successively with 100 ml each of CH_2Cl_2 , 6% DIPEA/ CH_2Cl_2 (two times), CH_2Cl_2 (two times). The resin was treated with p- [(R, S)- α -[1-(9H-fluoren-9-yl)-methoxyformamido]-2,4-dimethoxybenzyl]-phenoxyacetic acid (Fmoc-Linker) (7.01

g, 13.0 mmole), N-hydroxybenzotriazole (2.16 g, 16.0 mmole), and diisopropylcarbodiimide (2.04 ml, 13.0 mmol) in 100 mL 25% DMF/CH₂Cl₂ for 24 hours at room temperature. The resin was filtered and washed successively with 100 ml each of CH₂Cl₂ (two times), isopropanol (two times), DMF, and CH₂Cl₂ (three times). A Kaiser ninhydrin analysis was negative. The resin was dried under vacuum to yield 16.12 g of Fmoc-Linker-BHA resin. A portion of this resin (3.5 mg) was subjected to Fmoc deprotection and quantitative UV analysis indicated a loading of 0.56 mmol/g.

EXAMPLE 30

Preparation of Ac-Nle-Cyclo(Asp-Lys)-Asp-His-(D)Phe-Arg-Trp-Lys-NH₂



[0248] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Seven coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-His (Trt) (600 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Nle (430, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 1 mL acetic anhydride in 6% DIPEA/CH₂Cl₂ for


30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Ac-Heptapeptide resin. The Ac-heptapeptide resin was treated with 100 μL ethanedithiol, 100 μL dimethylsulfide, 250 μL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0249] 250 mg of the crude linear peptides was dissolved in 250 ml of DMF, 600 μL of N-methyl morpholine was added to have an apparent pH of 8.0. 300 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0250] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 60 mg (15 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₀H₆₉N₁₅O₉ cal: 1024 observed: m/z (1025 M+H).

EXAMPLE 31

Preparation of Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-Trp-Lys-NH₂



[0251] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Apc** (550 mg 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g of Pentyl-hexapeptide resin.


[0252] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification. 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was

added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0253] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 53 mg (13%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₂N₁₂O₈ cal: 1017 observed: m/z (1018 M+H).

EXAMPLE 32

Preparation of Penta-cyclo(Asp-Lys)-Asp-**Apc**-(D)Phe-Arg-(2)Nal-Lys-NH₂



[0254] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(2)Nal (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Apc** (550 mg 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and

HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0255] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0256] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0257] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₆H₇₃N₁₁O₈ cal: 1028 observed: m/z (1029 M+H).

EXAMPLE 33

Preparation of Penta-cyclo(Asp-Lys)-Asp-**Apc**-(D)Phe-Arg-N-methyl(2)Nal-Lys-NH₂



[0258] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Two coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(2)Nal (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol).

[0259] After Fmoc removal from 2-Nal residue, the resulting amine was converted to its 2-nitrobenzene sulfonyl derivative using 2-nitrobenzenesulfonyl chloride (5 eq., 426 mg, 1.93 mmol) and DIPEA (5 eq.) as the base in DMF. Washings were performed using DMF (6 x 30 ml) followed by CH₂Cl₂ (3 x 30 ml) and the resin was dried under vacuum. The sulfonamide obtained was subjected to methylation using triphenylphosphine (5 eq., 505 mg, 1.93 mmol), N, N-diethylazodicarboxylate (5 eq., 303 μ L, 1.93 mmol) and methanol (10 eq. 156 μ L, 3.85 mmol) in THF. Washings were performed using THF (6 x 30 ml) followed by CH₂Cl₂ (5 x 30 ml) and the resin was dried under vacuum. The 2-nitrobenzene sulfonyl group was then removed using 1,8-diazabicyclo [5.4.0] undec-7-ene (3 eq., 173 μ L, 1.16 mmol), 2-mercaptoethanol (5eq. 135 μ L, 1.93 mmol) in DMF. Washings were performed using DMF (3 x 30 ml), isopropanol (3 x 30 ml) followed by ethyl ether (3 x 30 ml) and the resin was dried under vacuum. The resulting N-Me-2-Nal residue was subjected to four coupling cycles, one cycle each with Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Apc** (550 mg 1.2 mmol) and HBTU (452 mg, 1.2

mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0260] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 235 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0261] 235 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0262] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 43 mg (10 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₇H₇₅N₁₁O₈ cal: 1042 observed: m/z (1043 M+H).

EXAMPLE 34

Preparation of Cyclo(succinic acid-Lys)-succinic acid-Apc-(D)Phe-Arg-Trp-Lys-NH₂



[0263] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Succinic anhydride** (600 mg, 6 mmol) in DMF with 1.1 ml of DIPEA.

[0264] The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0g Pentapeptide resin. The pentapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude 1 product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.


[0265] 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was

added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0266] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 40 mg (11%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₄₉H₆₃N₁₁O₇ cal: 918 observed: m/z (919 M+H).

EXAMPLE 35

Preparation of Cyclo(maleic acid-Lys)-maleic acid-Apc-(D)Phe-Arg-Trp-Lys-NH₂



[0267] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Maleic anhydride** (600 mg, 6 mmol) in DMF with the addition of HOBT (800 mg, 6 mmol), **without** DIPEA. The resin was filtered and


washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g Pentapeptide resin. The pentapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude product was dried under vacuum to yield 230 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0268] 230 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0269] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 38 mg (11%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₄₉H₆₁N₁₁O₇ cal: 916 observed: m/z (917 M+H).

EXAMPLE 36

Preparation of Cyclo(phthalic acid-Lys)-phthalic acid-**Apc**-(D)Phe-Arg-Trp-Lys-NH₂



[0270] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Phthalic anhydride** (660 mg, 6 mmol) in DMF with 1.1 ml of DIPEA.


[0271] The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0g Pentapeptide resin. The pentapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude 1 product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0272] 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0273] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 35 mg (10 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₃H₆₃N₁₁O₇ cal: 966 observed: m/z (967 M+H).

EXAMPLE 37

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-OHAp_c-(D)Phe-Arg-Trp-Lys-NH₂



[0274] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-OHAp_c** (565 mg

1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1g of Pentyl-hexapeptide resin.


[0275] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 225 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0276] 225 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0277] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (13%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₂N₁₂O₉ cal: 1033 observed: m/z (1034 M+H).

EXAMPLE 38

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-MeOApc-(D)Phe-Arg-Trp-Lys-NH₂



[0278] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-MeOApc** (600mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0279] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 235 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0280] 235 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0281] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 49 mg (12%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₉ cal: 1047 observed: m/z (1048 M+H).

EXAMPLE 39

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-EtOApc-(D)Phe-Arg-Trp-Lys-NH₂



[0282] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-EtOApc** (640mg

1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0283] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA, and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 235 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0284] 235 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0285] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 60 mg (14%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₆H₇₆N₁₂O₉ cal: 1061 observed: m/z (1062 M+H).

EXAMPLE 40

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-iPrOApc-(D)Phe-Arg-Trp-Lys-NH₂



[0286] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-iPrOApc** (660mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0287] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 260 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0288] 260 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0289] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 63 mg (15%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₇H₇₈N₁₂O₉ cal: 1075 observed: m/z (1076 M+H).

EXAMPLE 41

Preparation of Penta-cyclo(Asp-Lys)-Asp-3-MeOApc-(D)Phe-Arg-Trp-Lys-NH₂



[0290] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-

(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-3-MeOApc** (600 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0291] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 235 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0292] 235 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0293] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 49 mg (12%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₉ cal: 1047 observed: m/z (1048 M+H).

EXAMPLE 42

Preparation of Penta-cyclo(Asp-Lys)-Asp-**4-ClApc**-(D)Phe-Arg-Trp-Lys-NH₂



[0294] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-ClApc** (560 mg 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g of Pentyl-hexapeptide resin.


[0295] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 230 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0296] 230 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0297] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 49 mg (12 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₁N₁₂O₈Cl ; cal: 1051 observed: m/z (1052 M+H).

EXAMPLE 43

Preparation of Penta-cyclo(Asp-Lys)-Asp-4- MeApc-(D)Phe-Arg-Trp-Lys-NH₂



[0298] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2

mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-MeApc** (590mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0299] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0300] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0301] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₈ cal: 1031 observed: m/z (1032 M+H).

EXAMPLE 44

Preparation of Penta-cyclo(Glu-Lys)-Glu-Apc-(D)Phe-Arg-Trp-Lys-NH₂



[0302] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Fmoc-Glu (OBut)** (510 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.

[0303] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 255mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0304] 255 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0305] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 60 mg (15%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₈ cal: 1031 observed: m/z (1032M+H).

EXAMPLE 45

Preparation of Penta-cyclo(Asp-Orn)-Asp-Apc-(D)Phe-Arg-Trp-Orn-NH₂



[0306] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling

cycles were performed of one cycle each with **Fmoc-Orn (Boc)** (550 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.15 g of Pentyl-hexapeptide resin.


[0307] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0308] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuo, and the resulting reaction mixture was purified by HPLC.

[0309] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 53 mg (13%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₃H₇₀N₁₂O₈ cal: 1003 observed: m/z (1004 M+H).

EXAMPLE 46

Preparation of Penta-cyclo(Asp-Dbr)-Asp-**Apc**-(D)Phe-Arg-Trp-Dbr-NH₂



[0310] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with **Fmoc-Dbr (Boc)** (540 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.10 g of Pentyl-hexapeptide resin.

[0311] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted.


The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 220mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0312] 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0313] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 35 mg (9 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₈N₁₂O₈ cal: 989 observed: m/z (990 M+H).

EXAMPLE 47

Preparation of Penta-cyclo(Asp-Dpr)-Asp-Apc-(D)Phe-Arg-Trp-Dpr-NH₂



[0314] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling

cycles were performed of one cycle each with **Fmoc-Dpr (Boc)** (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g of Pentyl-hexapeptide resin.


[0315] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 200mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0316] 200 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0317] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 30 mg (8%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₁H₆₆N₁₂O₈ cal: 975 observed: m/z (976 M+H).

EXAMPLE 48

Preparation of Ac-cyclo(Asp-Dpr)-Asp-Apc-(D)Phe-Arg-Trp-Dpr-NH₂



[0318] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with **Fmoc-Dpr (Boc)** (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL acetic anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Acetyl-hexapeptide resin.

[0319] The Acetyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with \sim 2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 200mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0320] 210 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0321] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 28 mg (8%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₄₈H₆₀N₁₂O₈ cal: 933 observed: m/z (934 M+H).

EXAMPLE 49

Preparation of Cyclo(phthalic acid-Dpr)-phthalic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂



[0322] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling

cycles were performed of one cycle each with Fmoc-Dpr (Boc) (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Phthalic anhydride** (660 mg, 6 mmol) in DMF with 1.1 ml of DIPEA. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g Pentapeptide resin.


[0323] The pentapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0324] 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0325] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 30 mg (8 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₀H₅₇N₁₁O₇ cal: 924 observed: m/z (925 M+H).

EXAMPLE 50

Preparation of Cyclo(succinic acid-Dpr)-succinic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂



[0326] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Dpr (Boc) (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Succinic anhydride** (600 mg, 6 mmol) in DMF with 1.1 ml of DIPEA. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g Pentapeptide resin.


[0327] The pentapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude product was dried under vacuum to yield 220 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0328] 220 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0329] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 31 mg (8 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₄₆H₅₇N₁₁O₇ cal: 876 observed: m/z (877 M+H).

EXAMPLE 51

Preparation of Cyclo(maleic acid-Dpr)-maleic acid-Apc-(D)Phe-Arg-Trp-Dpr-NH₂



[0330] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Dpr (Boc) (530 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-

(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), **Maleic anhydride** (600 mg, 6 mmol) in DMF with the addition of HOBT (800 mg, 6 mmol), **without** DIPEA. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g Pentapeptide resin.


[0331] The pentapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude product was dried under vacuum to yield 230 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0332] 230 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0333] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 28 mg (8 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₄₆H₅₅N₁₁O₇ cal: 874 observed: m/z (875 M+H).

EXAMPLE 52

Preparation of Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Cit-Trp-Lys-NH₂



[0334] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Cit (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.3 g of Pentyl-hexapeptide resin.


[0335] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 300 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0336] 300 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0337] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 80 mg (20%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₁N₁₁O₉ cal: 1018 observed: m/z (1019 M+H).

EXAMPLE 53

Preparation of Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Ala-Trp-Lys-NH₂



[0338] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Ala (380 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and

HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.4 g of Pentyl-hexapeptide resin.


[0339] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 330 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0340] 330 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0341] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 87 mg (20%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₁H₆₅N₉O₈ cal: 932 observed: m/z (933 M+H).


EXAMPLE 54

Preparation of Ac-Nle-cyclo(Cys-Cys)-Cys-Apc-(D)Phe-Arg-Trp-Cys-NH₂



[0342] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Seven coupling cycles were performed of one cycle each with Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Apc** (550 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Nle (430, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 1 mL acetic anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Ac-Heptapeptide resin.

[0343] The Ac-heptapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted.

[illegible][illegible][illegible][illegible][illegible]

[0346] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-(D,L)-Atc** (510 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.15 g of Pentyl-hexapeptide resin.

[0347] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.


[0348] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0349] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main

peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray $C_{52}H_{68}N_{12}O_8$ cal: 989 observed: m/z (990 $M+H$).

EXAMPLE 56

Preparation of Penta-(Asp-Lys)-Asp-5-BrAtc-(D)Phe-Arg-Trp-Lys-NH₂ (peak 1)



[0350] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Br-(D,L)Atc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu^t) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0351] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 240 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0352] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0353] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The **first main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 26 mg (6 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₇N₁₂O₈Br cal: 1068 observed: m/z (1069 M+H).


EXAMPLE 57

Preparation of Penta-(Asp-Lys)-Asp-**5-BrAtc**-(D)Phe-Arg-Trp-Lys-NH₂ (peak 2)



[0354] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Br-(D,L)Atc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.

[0355] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 240 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[illegible][illegible][illegible][illegible][illegible]

mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Cl(D,L)Atc** (560 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0359] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0360] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0361] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. **The first main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 24 mg (6 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₇N₁₂O₈Cl cal: 1024 observed: m/z (1025 M+H).

EXAMPLE 59

Preparation of Penta-(Asp-Lys)-Asp-**5-ClAtc**-(D)Phe-Arg-Trp-Lys-NH₂ (peak 2)



[0362] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Cl-(D,L)Atc** (560 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin. The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was


dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0363] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0364] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. **The second main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 20 mg (4 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₇N₁₂O₈Cl cal: 1024 observed: m/z (1025 M+H).

EXAMPLE 60

Preparation of Penta-(Asp-Lys)-Asp-5-MeO-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂



[0365] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed

using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-MeO-(D,L)Atc** (600 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0366] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.


[0367] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0368] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized

to yield 55 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray $C_{53}H_{70}N_{12}O_9$ cal: 1019 observed: m/z (1020 $M+H$).

EXAMPLE 61

Preparation of Penta-(Asp-Lys)-Asp-**5-EtO-(D,L)Atc**-(D)Phe-Arg-Trp-Lys-NH₂



[0369] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-EtO-(D,L)Atc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.3 g of Pentyl-hexapeptide resin.


[0370] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 260 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0371] 260 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0372] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 58 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₂N₁₂O₉ cal: 1033 observed: m/z (1034 M+H).

EXAMPLE 62

Preparation of Penta-(Asp-Lys)-Asp-5-iPrO-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂



[0373] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-iPrO-(D,L)Atc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0374] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0375] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0376] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 58 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₉ cal: 1047 observed: m/z (1048 M+H).

EXAMPLE 63

Preparation of Penta-(Asp-Lys)-Asp-5-Me-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂



[0377] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed

using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Me-(D,L)Atc** (590 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0378] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 260 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.


[0379] 260 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0380] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized

to yield 62 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray $C_{53}H_{70}N_{12}O_8$ cal: 1003 observed: m/z (1004 $M+H$).

EXAMPLE 64

Preparation of Penta-(Asp-Lys)-Asp-5-Et-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂



[0381] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Et-(D,L)Atc** (600 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.3 g of Pentyl-hexapeptide resin.

[0382] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for


180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0383] 245 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0384] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (12 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₄H₇₂N₁₂O₈ cal: 1017 observed: m/z (1018 M+H).

EXAMPLE 65

Preparation of Penta-(Asp-Lys)-Asp-5-iPr-(D,L)Atc-(D)Phe-Arg-Trp-Lys-NH₂



[0385] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-iPr-(D,L)Atc** (600 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0386] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.


[0387] 245 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0388] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main

peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 54 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray $C_{55}H_{74}N_{12}O_8$ cal: 1031 observed: m/z (1032 $M+H$).

EXAMPLE 66

Preparation of Penta-(Asp-Lys)-Asp-5-BrAtc-(D)Phe-Cit-Trp-Lys-NH₂ (peak 1)



[0389] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Cit (480mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-5-Br-(D,L)Atc (620 mg 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0390] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 240 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 260 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0391] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0392] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The **first main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 24 mg (5,6 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₆N₁₁O₉Br cal: 1069 observed: m/z (1070 M+H).

EXAMPLE 67

Preparation of Penta-(Asp-Lys)-Asp-**5-BrAtc**-(D)Phe-Cit-Trp-Lys-NH₂ (peak 2)



[0393] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Cit** (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Br-(D,L)Atc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0394] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 240 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0395] 240 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0396] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The **second main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 22 mg (4.8 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₆N₁₁O₉Br cal: 1069 observed: m/z (1070 M+H).

EXAMPLE 68

Preparation of Penta-(Asp-Lys)-Asp-5-ClAtc-(D)Phe-Cit-Trp-Lys-NH₂ (peak 1)



[0397] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2

mmol), **Fmoc-Cit** (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-Cl-(D,L)Atc** (560 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0398] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0399] 245 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0400] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. **The first main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 22 mg (5.8 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₆N₁₁O₉Cl cal: 1024 observed: m/z (1025 M+H).

EXAMPLE 69

Preparation of Penta-(Asp-Lys)-Asp-5-ClAtc-(D)Phe-Cit-Trp-Lys-NH₂ (peak 2)



[0401] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Cit (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-5-Cl-(D,L)Atc (560 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0402] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0403] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0404] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. **The second main peak** was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 20 mg (5.4 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₆₆N₁₁O₉Cl cal: 1024 observed: m/z (1025 M+H).

EXAMPLE 70

Preparation of Ac-Nle-cyclo(Cys-Cys)-(D,L)Atc-(D)Phe-Arg-Trp-Cys-NH₂



[0405] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed

using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Seven coupling cycles were performed of one cycle each with Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-(D,L)Atc** (550 mg, 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Nle (430 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 1 mL acetic anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g of Ac-Heptapeptide resin.

[0406] The Acetyl-hepta peptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA, and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid.


[0407] This crude linear peptide was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg of purified linear peptide.

[0408] The purified linear peptide was dissolved in 2 ml of DMSO, diluted with 500 ml of water and the pH was adjusted to pH 8.0 with NH₄OH. O₂ was bubbled into the solution and the cyclization was monitored by HPLC. Typically the cyclization was completed within 24-48 hours. The solution was lyophilized and the material dissolved in

CH₃COOH and was subjected to preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 20 mg (5.0 %) of purified cyclic peptide. This compound was homogeneous by HPLC. LR-Electrospray C₅₁H₆₆N₁₂O₈S₂ cal: 1039 observed: m/z (1040 M+H).

EXAMPLE 71

Preparation of Penta-cyclo(Cys-Cys)-Cys-**5-Br(D,L)Atc**-(D)Phe-Arg-Trp-Cys-NH₂



[0409] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Seven coupling cycles were performed of one cycle each with Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-5-BrAtc** (620 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Cys (Trt) (710 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 1 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50

ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Ac-Heptapeptide resin.


[0410] The pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA, and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 240 mg of an off-white solid.

[0411] This crude linear peptide was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 50 mg of purified linear peptide.

[0412] The purified linear peptide was dissolved in 2 ml of DMSO, diluted with 500 ml of water and the pH was adjusted to pH 8.0 with NH₄OH. O₂ was bubbled into the solution and the cyclization was monitored by HPLC. Typically the cyclization was completed within 24-48 hours. The solution was lyophilized and the material dissolved in CH₃COOH and was subjected to preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 22 mg (5.2 %) of purified cyclic peptide. This compound was homogeneous by HPLC. LR-Electrospray C₄₈H₆₀N₁₁O₇S₂ Br cal: 1047 observed: m/z (1048 M+H).

EXAMPLE 72

Preparation of Penta-cyclo(Asp-Lys)-Asp-**Appc**-(D)Phe-Arg-Trp-Lys-NH₂



[0413] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Appe** (550mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0414] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0415] 245 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0416] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 57 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₃H₇₁N₁₃O₈ cal: 1018 observed: m/z (1019 M+H).

EXAMPLE 73

Preparation of Penta-cyclo(Asp-Lys)-Asp-2-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂



[0417] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling

cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-2-MeAppc** (570mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0418] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0419] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0420] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 61 mg (15 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₄H₇₃N₁₃O₈ cal: 1032 observed: m/z (1033M+H).

EXAMPLE 74

Preparation of Penta-cyclo(Asp-Lys)-Asp-2-iPrAppc-(D)Phe-Arg-Trp-Lys-NH₂



[0421] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-2-iPrAppc** (600 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0422] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the


crude linear product was dried under vacuum to yield 245 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0423] 245 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0424] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 52 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₆H₇₇N₁₃O₈ cal: 1060 observed: m/z (1061 M+H).

EXAMPLE 75

Preparation of Penta-cyclo(Asp-Lys)-Asp-3-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂



[0425] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling

cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-3-MeAppc** (570mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0426] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 248 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0427] 248 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0428] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₄H₇₃N₁₃O₈ cal: 1032 observed: m/z (1033 M+H).

EXAMPLE 6

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-MeAppc-(D)Phe-Arg-Trp-Lys-NH₂



[0429] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-MeAppc** (570mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.

[0430] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted.


The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 254 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0431] 254 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0432] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 57 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₄H₇₃N₁₃O₈ cal: 1032 observed: m/z (1033 M+H).

EXAMPLE 77

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-ClAppc-(D)Phe-Arg-Trp-Lys-NH₂



[illegible]

Figure 6. The effect of the initial concentration of the monomer (C_0) on the polymerization rate at different temperatures. The reaction conditions were: $[AIBN] = 0.005 \text{ mol/L}$, $[M] = 0.05 \text{ mol/L}$, $[K_2S_2O_8] = 0.005 \text{ mol/L}$, $[NaHCO_3] = 0.005 \text{ mol/L}$, $[NaHSO_3] = 0.005 \text{ mol/L}$, $[Na_2SO_3] = 0.005 \text{ mol/L}$, $[Na_2S_2O_4] = 0.005 \text{ mol/L}$, $[Na_2S_2O_5] = 0.005 \text{ mol/L}$, $[Na_2S_2O_8] = 0.005 \text{ mol/L}$, $[Na_2S_2O_7] = 0.005 \text{ mol/L}$.


Figure 6. The effect of the initial concentration of the monomer (C_0) on the polymerization rate at different temperatures. The reaction conditions were: $[AIBN] = 0.005 \text{ mol/L}$, $[M] = 0.05 \text{ mol/L}$, $[K_2S_2O_8] = 0.005 \text{ mol/L}$, $[NaHCO_3] = 0.005 \text{ mol/L}$, $[NaHSO_3] = 0.005 \text{ mol/L}$, $[NaNO_2] = 0.005 \text{ mol/L}$, $[NaCl] = 0.005 \text{ mol/L}$, $[NaBr] = 0.005 \text{ mol/L}$, $[NaI] = 0.005 \text{ mol/L}$, $[NaOH] = 0.005 \text{ mol/L}$, $[Na_2SO_4] = 0.005 \text{ mol/L}$, $[Na_2CO_3] = 0.005 \text{ mol/L}$, $[Na_2SO_3] = 0.005 \text{ mol/L}$, $[Na_2S_2O_8] = 0.005 \text{ mol/L}$, $[Na_2S_2O_7] = 0.005 \text{ mol/L}$, $[Na_2S_2O_8] = 0.005 \text{ mol/L}$, $[Na_2S_2O_7] = 0.005 \text{ mol/L}$.

Figure 6. The effect of the initial concentration of the monomer (C_0) on the polymerization rate at different temperatures. The reaction conditions were: $[AIBN] = 0.005 \text{ mol/L}$, $[M] = 0.05 \text{ mol/L}$, $[K_2S_2O_8] = 0.005 \text{ mol/L}$, $[NaHCO_3] = 0.005 \text{ mol/L}$, $[NaHSO_3] = 0.005 \text{ mol/L}$, $[NaNO_2] = 0.005 \text{ mol/L}$, $[NaCl] = 0.005 \text{ mol/L}$, $[NaBr] = 0.005 \text{ mol/L}$, $[NaI] = 0.005 \text{ mol/L}$, $[NaOH] = 0.005 \text{ mol/L}$, $[Na_2SO_4] = 0.005 \text{ mol/L}$, $[Na_2CO_3] = 0.005 \text{ mol/L}$, $[Na_2SO_3] = 0.005 \text{ mol/L}$, $[Na_2S_2O_8] = 0.005 \text{ mol/L}$, $[Na_2S_2O_7] = 0.005 \text{ mol/L}$, $[Na_2S_2O_8] = 0.005 \text{ mol/L}$, $[Na_2S_2O_7] = 0.005 \text{ mol/L}$.

peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 55 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: $C_{53}H_{70}N_{13}O_8Cl$ cal: 1032 observed: m/z (1033 $M+H$).

EXAMPLE 78

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-**PhOAppc**-(D)Phe-Arg-Trp-Lys-NH₂



[0437] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-4-PhOAppc** (650 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0438] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 270 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0439] 270 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0440] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 58 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₉H₇₅N₁₃O₉ cal: 1110 observed: m/z (1111 M+H).

EXAMPLE 79

Preparation of Penta- (Asp-Lys)-Asp-3-MeO-Apoc-(D)Phe-Arg-Trp-Lys- NH₂



[0441] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-3-MeOAAppc** (580 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.2 g of Pentyl-hexapeptide resin.


[0442] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0443] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0444] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 54 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₄H₇₃N₁₃O₉ cal: 1048 observed: m/z (1049 M+H).

EXAMPLE 80

Preparation of Penta-cyclo(Asp-Lys)-Asp-4-Adpc-(D)Phe-Arg-Trp-Lys-NH₂



[0445] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Adpc** (620 mg 1.2

mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0446] The Pentyl-hexapeptide resin was treated with 100 µL ethanedithiol, 100 µL dimethylsulfide, 250 µL anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 242 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0447] 242 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0448] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 48 mg (11%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₆₀H₇₆N₁₂O₈ cal: 1093 observed: m/z (1094 M+H).

EXAMPLE 81

Preparation of Penta-cyclo(Asp-Lys)-Asp-Achc-(D)Phe-Arg-Trp-Lys-NH₂



[0449] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Ahc** (560 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBu) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.1 g of Pentyl-hexapeptide resin.


[0450] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 250 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0451] 250 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0452] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 52 mg (13 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray: C₅₄H₇₈N₁₂O₈ cal: 1023 observed: m/z (1024 M+H).

EXAMPLE 82

Preparation of Penta-cyclo(Asp-Lys)-Asp-**Abc**-(D) Phe-Arg-Trp-Lys-NH₂



[0453] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Trp (520 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-

(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-Abc** (530 mg 1.2 mmol) and HBTU (452 mg, 0.6 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 13 g of Pentyl-hexapeptide resin.


[0454] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL trifluoroacetic acid at room temperature for 180 min. The resin was filtered off, washed with ~2 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and recentrifuged and the crude linear product was dried under vacuum to yield 255 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0455] 255 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 μ L of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum and the resulting reaction mixture was purified by HPLC.

[0456] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 58 mg (14 %) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₂H₇₆N₁₂O₈ cal: 997 observed: m/z (998 M+H).

EXAMPLE 83

Preparation of Penta-cyclo(Asp-Lys)-Asp-Apc-(D)Phe-Arg-(2S,3S) **beta** methyl-Trp-Lys-NH₂



[0457] Fmoc-Linker-BHA resin (720 mg, 0.4 mmol) from Example 29 were subjected to solid phase synthesis using Protocol 1 described above. All couplings were performed using HBTU in DMF as the coupling agent and DIPEA (3 equiv.) as base. Six coupling cycles were performed of one cycle each with Fmoc-Lys (Boc) (565 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), **Fmoc-(2S,3S) beta methyl (nMes)Trp** (616 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Arg (Pmc) (800 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-(D)Phe (480 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Apc (550 mg 1.2 mmol) and HBTU (452 mg, 1.2 mmol), Fmoc-Asp (OBut) (500 mg, 1.2 mmol) and HBTU (452 mg, 1.2 mmol). The peptide resin was carried through steps 1 - 5 of protocol 1, washed with CH₂Cl₂ (three times) and treated with 2 mL valeric anhydride in 6% DIPEA/CH₂Cl₂ for 30 minutes. The resin was filtered and washed successively with 50 ml each of CH₂Cl₂ (two times), isopropanol, and CH₂Cl₂ (three times). The resin was dried under vacuum to yield 1.0 g of Pentyl-hexapeptide resin.

[0458] The Pentyl-hexapeptide resin was treated with 100 μ L ethanedithiol, 100 μ L dimethylsulfide, 250 μ L anisole, and 10 mL HF at 0C for 60 min. HF was evaporated, and the resin was washed with ethylacetate filtered off, washed with ~5 ml TFA and the filtrates precipitated in chilled ethyl ether. The precipitates were centrifuged and the ether layer decanted. The residue was washed with two or three volumes of Et₂O and

recentrifuged and the crude linear product was dried under vacuum to yield 180 mg of an off-white solid. The crude peptide was subjected to cyclization without purification.

[0459] 180 mg of the crude linear peptides was dissolved in 220 ml of DMF, 500 µL of N-methyl morpholine was added to have an apparent pH of 8.0. 280 mg of BOP was added and the cyclization was monitored by HPLC. Typically the cyclization was completed within 18-24 hours. 10 ml of water was added to stop the reaction, DMF was evaporated in vacuum, and the resulting reaction mixture was purified by HPLC.

[0460] This crude cyclic material was purified by preparative HPLC on a Vydac C18-column (2.5 x 20 cm) and eluted with a linear gradient of 20-60% B (buffer A: 0.1% TFA/H₂O, buffer B: 0.1% TFA/CH₃CN) in 90 min., flow rate 8 ml/min, detection 280 nm. The main peak was cut by analytical HPLC analysis of collected fractions, pooled and lyophilized to yield 40 mg (10%) of a white, amorphous powder. This compound was homogeneous by HPLC. LR-Electrospray C₅₅H₇₄N₁₂O₈ cal: 1031 observed: m/z (1032 M+H).

BIOLOGICAL ACTIVITY EXAMPLE:

Example A: Agonist Assay

Method

[0461] Description: HEK 293 cells transfected with either the MC-4 receptor or MC-1 receptor were grown in 96 well plates. The cells were stimulated with either 100nM NDP-αMSH or screening compounds. Cyclic AMP was extracted from the cells and concentrations were determined using a Biotrak-cAMP SPA assay. Agonists were identified as those compounds causing an increase in cAMP.

[0462] Cell Culture: HEK 293 cells transfected with either the MC-4 receptor or MC-1 receptor were cultured in 75cm² flasks in D-MEM supplemented with 10% FCS and 500µg/ml G418. Cells were trypsinized and split 1:3 into 96 well flat-bottom tissue culture treated plates. Cells were stimulated at confluence (day 2-4).

[0463] cAMP Response: Compounds serially diluted in 100% DMSO were further diluted 1:200 (2.5µl compound dilution + 500µl media) in D-MEM containing 10%FBS and 0.1mM IBMX. For unstimulated cells, 2.5µl of DMSO was added to 500µl of media. For NDP-αMSH stimulated cells, 2.5µl of 20µM NDP-αMSH in 100% DMSO was added to 500µl of media (final conc. 100nM). Final concentration of DMSO in all wells was 0.5%.

[0464] Note: Each sample was run in duplicate on separate plates.

[0465] Culture medium was removed from confluent 96 well culture plates and replaced with 200µl of above dilutions into the appropriate wells. The plates were incubated for 1hr at RT. The media was removed, and the plates were washed 1x with 200µl well of PBS. CAMP was extracted by the addition of 60µl 70% ethanol (stored in the refrigerator). After a 30min extraction period, 10µl ethanol extract was transferred to the cAMP assay plate or samples were stored at -20°C until the cAMP assay was performed.

[0466] cAMP Assay: The extracted samples and all reagents included in the kit were brought to room temperature. To a 96 well OptiPlate, 10µl ethanol extract, 40µl assay buffer, 50ul [125I]cAMP, 50µl antiserum and 50µl SPA beads were added. The total well volume after addition was 200µl. The plates were sealed and incubated for 15-20 hr at room temperature. [125I]cAMP binding to the SPA beads was determined by counting each plate for 2 minutes on a Packard TopCount™.

[0467] Note: Each plate contained samples of controls for unstimulated cells and NDP-αMSH for stimulated cells.